THE PLANT DISEASE REPORTER

Issued by

CROPS RESEARCH DIVISION

INDEX TO VOLUME 45

Prepared by Nellie W. Nance and Mary K. Burk

Crops Research Division
Volume 45

Plant Industry Station, Beltsville, Maryland
Index Number

AUTHOR INDEX

AICHELE, MURIT D., (45)
ALBERT, J. J., 759
ALCORN, STANLEY M., 587
ALEXANDER M., (487)
ALFORD, D. M., 856
ALLAN, R. E., 778
ANCALMO, OSCAR, 281, 576, 666, 826, 900
APPLE, J. L., (362), 968
APT, WALTER J., 290
ARK, P. A., (85)
ARNY, D. C., (192)
ASAI, G. N., (907)
ASHRI, AMRAM, 146, 153
ASHWORTH, LEE J., Jr., 220
ATLEE, C. B., Jr., (739)
AVERRE, C. W., III, (435)
AYCOCK, ROBERT, 620
AYOUB, SADEK M., 940

BAIN, DOUGLAS C., 814
BAJAJ, B. S., 260
BAKER, RALPH, 76
BALDACCI, ELIO, (912)
BARNES, GEORGE L., 142, 426
BARTLETT, ALLAN, (628)
BATJER, L. P., 255
BEAN, GEORGE A., 901
BENSON, N. R., (255)
BENTAL, A., (356)
BERKENKAMP, BILL B., 826
BERNARD, R. L., (789)
BERRY, CHARLES R., 152
BERRY, FREDERICK H., 167, 576
BILBRUCK, JAMES D., 128
BIRCHFIELD, W., 671
BLANK, LESTER M., 241
BLODGETT, EARLE C., 45, (258), (912)
BOEWE, G. H., 393
BOHN, G. W., 232, (677)
BORDERS, HUEY I., 972
BOSHER, J. E., (481)
BOWMAN, D. H., (764)
BOWMAN, JACOB N., (632)
BOYCE, JOHN S., Jr., (306), 386
BRAVERMAN, SAMUEL W., 199
BREECE, JAMES R., (796)
BRETT, CHARLES H., (330)
BRIGGLE, L. W., 846
BRIGHAM, RAYMOND D., 124
BRINKERHOFF, L. A., 126
BROWDER, L. E., 894
BRUEHL, G. W., (803), 884
BURBUTIS, P. P., (120)
BURK, L. G., 734
BURNETT, HARRY C., 697
BUSHONG, J. W., 100

CALDWELL, R. M., (444)
CALPOUZOS, L., (72)
CANNON, ORSON S., (354)
CAPPELLINI, R. A., 301
CAROSELLI, NESTOR E., 24, (965)
CATION, DONALD, 109
CHAPMAN, R. A., (140)
CHIARAPPA, LUIGI, (912)
CHRISTENSEN, J. J., (448)
CHUNG, BONG KOO, (680)
CHUNG, HOO SUP, 680
CLARK, W. A., (384), (755)
COBB, F. W., Jr., (736)
COHN, E., 505
COLE, JOHN R., 911
COMMITTEE of North American Wheat Leaf Rust Research Workers, 444
COMPTON, O. C., (634)
CONKLIN, J. G., (899)
CONVERSE, RICHARD H., 882
COOPER, W. E., (173), (219), 674
CORDY, C. B., 891
COTTER, R. U., (448)
COUCH, HOUSTON B., (616)
CRAIGMILES, J. P., (216)
CRITTENDEN, H. W., 190
CROSSAN, D. F., 120, (874)
CRUM, R. A., (544)
CURL, E. A., 517

DAMSTEEGT, V. D., (884)
DARBY, J. F., 58, 552
DAS, G. N., 979
DAVIDSON, ROSS W., 836
DAVIS, CLAUDE J., (470)
DAVIS, R. A., (544)
DAVIS, WILLIAM C., (281)
DEBA, RAPHAEL, (878)
DEEP, IRA W., 628

DEGMAN, E. S., (255)
DELL, T. R., (38), (352), (660)
DESAI, M. V., 203
DEWEY, WADE G., 663
DeWOLFE, T. A., (264), (268)
DeZEEUW, D. J., 544, (605)
DIAB, KHALIL A., (932)
Di EDWARDO, A. A., 67
DOOLING, O. J., (576), 749
DRAKE, CHARLES R., 572
DRAKE, R. J., (682)
DRIVER, CHARLES H., 38, 352, 660
DROPKIN, V. H., (734)
DuCHARME, E. P., (62), (564), 742
DUKES, P. D., 362
DUNN, STUART, (574)
DURAN, RUBEN, 475
DURBIN, RICHARD D., (260)

EDGINGTON, L. V., (29)
EDWARDSON, JOHN R., (725), 958
EGLITIS, MAKSIS, (296)
ELGINDI, DAWOOD M., (928) (930)
ELLIOTT, EDWARD S., 470
ENDO, R. M., 869
EPSTEIN, ABRAHAM H., 826
ESCOBAR, PEDRO, (119)
EVERETTE, GEORGE A., (140)

FELDMAN, A. W., (62), 564, (782)
FERGUS, CHARLES L., 736, 916
FERRIS, VIRGINIA R., 789
FIELDHOUSE, D. J., (120)
FIGARO, PEGGYBETH, (730)
FORBES, I. L., 274
FORBES, IAN Jr., (725), (958)
FORDYCE, CLAUDE, Jr., (30), (108)
FRAZIER, N. W., 649
FRENCH, D. W., (749)
FRIDLUND, P. R., 687
FULTON, ROBERT H., (368)
FUMAGALLI, ASTOLFO, (227), (788)
FUTRELL, MAURICE C., (220)

GALVEZ, GUILLERMO E., 949, 954
GARDNER, W. S., (720)
GENTILE, A. G., (562)
GIBLER, JOHN W., 334
GILBERTSON, R. L., (389)
GILL, D. L., 300
GILMER, R. M., 228, 608, 612
GINSBURG, O., (945)
GOENAGA, ALVARO, (756)
GOHEEN, A. C., 641
GOLD, A. H., (649)
GOLDEN, A. MORGAN, (26)
GOOD, J. M., 46, 976
GOODE, M. J., 690
GORDON, PHILIP N., (756)
GOSS, ROY L., (112), (296)
GOULD, CHARLES J., 112, (290), 296, 508
GRAHAM, S. O., 41
GRAHAM, T. W., 692
GRANT, T. J., 416
GREELEY, LESTER W., (755)
GREEN, H. J., (134)
GREEN, RALPH J., Jr., 30, (108), 288, 696
GREENE, GEORGE L., 390
GUBA, EMIL F., 102
GUTHRIE, F. E., (225)

HAAS, J. H., (858)
HAASIS, FRANK A., 491
HABECK, DALE H., 330
HADWIGER, LEE A., 373
HAFSTAD, GEORGE E., 152, 681
HALL, CHARLES V., (373)
HALPIN, JAMES E., 298
HAMNER, C. L., (605)
HANKS, R. W., (742)
HANNA, G. C., 562
HARMON, D. L., (384)
HART, W. H., (739)
HARTMANN, GEORGE C., 938
HEDDEN, ORVE K., (380)
HELTON, A. W., 500, 591, 918, 920
HENDERSON, WARREN R., 272
HENNEBERRY, T. J., (652)
HENRY, S. E., (95)
HEPTING, GEORGE H., 889
HERR, ROBERT R., 399
HEUBERGER, J. W., (759)
HIKIDA, H. R., 388
HILL, A. C., (720)
HIMELICK, E. B., 180, (351)
HINE, RICHARD B., 462, 868
HODGES, CHARLES S., 134, 745
HOLMES, FRANCIS W., 74
HOLTON, C. S., (5)
HOOKER, A. L., 236, 780
HOPPE, PAUL E., 99, 787
HORN, N. L., 818
HOSFORD, ROBERT M., Jr., 482
HOWARD, CHARLES M., (470)
HUGUELET, J. E., 368
HUNTER, R. E., (126)
HUSAIN, S. M., 263

INMAN, ROBERT, (799)

JACKSON, CURTIS R., 512, 798
JENKINS, ANNA E., 145
JENNINGS, PETER R., (334), (949)
JEWELL, F. F., 639
JOHNSON, E. M., 140
JOHNSON, HOWARD W., 542
JOHNSTON, C. O., (444), 446, 661, (894)
JOHNSTON, T. H., (95)
JONES, BRIAN M., 366, 794
JONES, J. E. (671)
JONES, JOHN PAUL, 168, 376, (542)
JONES, THOMAS W., 506, 980

KARAS, J. G., 605
KEANE, F. W. L., (204)
KEIL, HARRY L., 10
KENDRICK, E. L., 5
KENKNIGHT, GLENN, 304
KERR, THOMAS, (550)
KIESLING, RICHARD L., (698), (941)
KILPATRICK, R. A., 286, 341, 717, 899, 926
KIMBLE, K. A., (562)
KINGSLAND, GRAYDON C., 845
KLINE, D. M., (644)
KLINGNER, A., 235
KLOTZ, L. J., 264, 268
KRAMER, C. L., 309
KREITLOW, K. W., (717)
KRULL, CHARLES F., (334)
KULIK, M. M., 907
KWACK, BEYOUNG H., 574

LAIRD, E. F., Jr., 877
LANGDON, KENNETH R., 248
LASSMANN K., DOROTHEA, 981
LATHAM, A. J., 866
LEAPHART, CHARLES D., (722)
LEAR, BERT, 739
LINN, M. B., (866)
LLOYD, A. B., 422
LOCKWOOD, JOHN L., (422), 569
LOEGERING, W. Q., 384, 444
LOOS, CLIVE A., 457
LÓPEZ, ARMANDO ALAS, (576)
LOTT, T. B., 204
LUCAS, G. B., 159
LUEPSCHEN, N. S., 557
LUESCHOW, WILLIAM C., (152)
LUTTRELL, E. S., 216
LYDA, STUART D., (636)
LYON, HOWARD, (312)

MacSWAN, I. C., (891)
MADISON, JOHN H., 892
MAHADEVAN, A., 965
MAI, W. F., 151, 211
MAIER, CHARLES R., 276, 804, 808, 960
MALO, SIMON E., 20
MANKAU, R., 164
MARSH, PAUL B., 550
MARTINEZ, A. P., 560
MARTINSON, CHARLIE, (76)
MAY, CURTIS, 78, 777
McALPINE, ROBERT G., 196
McCAIN, ARTHUR H., 151
McCARTER, STATES M., (298)
McCLELLAN, W. D., (508)
McDANIEL, M. C., (900)
McFADDEN, LORNE A., 16
McKINNEY, H. E., (497)
McKINNEY, H. H., (384), 755
McLEAN, D. M., 137, 728
MEYER, H. M., (137)
MICHEL, L. J., (974)
MIELKE, JAMES L., (831)
MILBRATH, J. A., 520
MILLER, JAMES D., 657
MILLER, P. M., 42
MILLER, PAUL R., 79, 236, 394, 750
MILLER, V. L., (112), (508)
MINZ, G., (505)
MITCHELL, R., 487
MOLNAR, A. C., 854
MONTGILLION, M. D., (244), (933)
MOORE, BOBBY, (530)
MOORE, LAURENCE D., 616
MOREIRA, SYLVIO, (416)
MORRISON, LOU S., (441)
MORTON, DONALD J., 348
MOTSINGER, R. E., 335
MURRAY, B. R., (731)
MUSE, D. P., (858), (863)

NAYUDU, M. V., (29)
NEELY, DAN, (180), 351
NELSON, R. R., 644
NICHOLS, LESTER P., 344, 346
NICKELL, LOUIS G., 756
NIELSEN, L. W., (330)
NIEMCZYK, H. D., 698
NORGREN, R. L., (746)
NORMAN, SHIRLEY M., (475)
NUSBAUM, C. J., 225, (491)

OGAWA, JOSEPH M., 636, (659)
ORCHARD, W. R., 308
ORLOB, G. B., 192, 466
OSBORNE, W. W., 812
OTEIFA, BAKIR A., 928, 930, 932

PADEN, JOHN W., (176)
PADY, S. M., (894)
PALMER, JOHN G., 652
PALTI, J., 31
PARRIS, G. K., 530
PARTRIDGE, A. D., (506)
PATEL, K. P., (203)
PATTON, ROBERT F. (836)
PÉREZ, J. ENRIQUE, 481
PETERSEN, D. H., 186
PETERSON, GLENN W., (799)
PETERSON, JOSEPH L., 208
PETERSON, ROGER S., 472
PHILLIPS, DOUGLAS J., (76)
PIRONE, T. P., (274), (856)
PONTIS-VIDELA, RAFAEL E., (235)
POWELL, DWIGHT, (100)
POWERS, H. R., Jr., 306
PRIDHAM, T. G., (244)
PROTACIO, D. B., 824
PROVVIDENTI, R., (160)
PURDY, LAURENCE H., (522)

RABB, R. L., (225)
RADEWALD, J. D., (411)

RANIERE, L. C., 844
RAWLINS, T. E., 598
REFATTI, ELVIO, (912)
REICHERT, I., 356, 945
REINKING, OTTO A., 411, 599
RESCONICH, E. C., (85)
REYES, TIBURCIO T., 185
REYNOLDS, J. E., (520)
REYNOLDS, JOHN F. (681)
RHOADES, H. L., 54
RICH, AVERY E., (128), (574), (845), (899)
RIGGS, R. D., 392
ROBERTS, A. N., 634
RODRIGUEZ, RICARDO A., 980
ROGERS, N. F., (576)
ROHDE, C. R., 522
ROTHMAN, P. G., 764
RUBIN, HERBERT L., 375

SAKIMURA, K., 766, 772
SALIBE, ARY A., (416)
SALISBURY, P. J., 481
SAMBORSKI, D. J., (444)
SAMSON, R. W., (30), 539
SASSER, J. N., 173, (620)
SCHAFER, J. F., (444)
SCHAREN, A. L., (846)
SCHENCK, N. C., 841
SCHIEBER, EUGENIO, 119, 227, 425, 788, 981
SCHINDLER, A. F., 413, 747
SCHLEGEL, D. E., (85)
SCHMIDT, R. A., (736)
SCHNATHORST, W. C., (641)
SCHREIBER, LAWRENCE R., 108
SCHROEDER, D. B., 749
SCHROEDER, W. T., 160
SCHWARTZ, RICHARD, (901)
SCIARONI, R. H., (739)
SEN GUPTA, P. K., (979)
SEQUEIRA, LUIS, 435
SEYMOUR, C. P., (900)
SHALLA, THOMAS A., 912
SIGURBJORNSSON, BJORN, (211)
SILBER, GUSTAVE, 310
SILIÉZAR, GUILLERMO RAMOS, 900
SILVERBORG, S. B., 389
SIMONS, M. D., 974
SINCLAIR, J. B., 625, (856)
SITTERLY, WAYNE R., 200
SKOTLAND, C. B., 51
SLEETH, BAILEY, (728)
SMALE, B. C., 244, 933
SMITH, A. L., (219)
SMITH, HARLAN E., 219
SMITH, K. M., (85)
SMITH, RALPH E., 632, 700, 702, 705, 709, 712
SMITH, T. E., 548, 549
SOSA, O. NERY, (119)
SOTOMAYOR-RÍOS, A., (72)
SPRAGUE, RODERICK, 106, 730, 803
SPURR, HARVEY W., Jr., 941
STALEY, JOHN M., 312
STALL, ROBERT E., 12
STAMBAUGH, WILLIAM J. (736), (916)
STANTON, THOMAS H., 659
STESSEL, G. J., 26
STEVENSON, JOHN A., 666
STEWART, DONALD M., 448, (657)
STRETCH, A. W., (301)
STRONG, M. C., 392
STRUBLE, F. BEN, (248), 441
STUCKEY, IRENE H., 527
SUIT, R. F., 62, 454, (564), 782
SWARTWOUT, H. G., (366), (794)

TAMMEN, J., 858, 863
TAYLOR, D. P., 746
TEMPLETON, G. E., 95
THEIS, T., 72
THOMASON, IVAN J., 497, 577
THOMPSON, GEORGE E., (152), 731
THOMPSON, H. C., 839
THOMPSON, HUGH E., (309)
THURSTON, H. DAVID, (949), (954)
TICKNOR, R. L., (634)
TODD, EDWIN H., 178
TODD, FURNEY A., 319
TOOLE, E. H., (933)
TOOLE, E. RICHARD, 78
TOWNSLEY, W. W., Jr., (120), 874
TRAYLOR, JACK A., (796)
TRESHOW, MICHAEL, 354, 720
TUITE, JOHN, 212

UI, TADAO, (102)

VALLEAU, W. D., (140)
Van ADRICHEM, M. C. J., (308)
VanDENBURGH, ROBERT, (120)
VAN DOREN, ARCHIE, (730)
VAN GUNDY, S. D., (577)
VARNEY, EUGENE H., 393
VEENSTRA, M. A., (368)
VIÉGAS, AHMÉS P., (145)
VOGEL, O. A., (778)
VOLCANI, ZAFRIRA, 823

WAGENER, WILLIS W., 831
WAGNON, H. KEITH, 796
WALKER, E. K., 583
WALKER, J. C., 29
WALKER, J. T., (380)
WALLACE, JAMES M., 682
WALTON, G. S., (301)
WATSON, R. D., 176, 289
WEATHERS, L. G., (877)
WEBB, R. E., 677, 851
WEBER, DARRELL J., (636)
WEIHING, JOHN L., 799
WELLS, DARRELL G., 878
WELLS, HOMER D., 725, (958)
WELLS, J. C., (491)
WHITAKER, THOMAS W., (232)

WICKER, ED F., 722
WILLIAMS, HAROLD E., (796)
WILSON, CHARLES L., 96, 900, 957
WILSON, J. D., 282, 380, 534
WILSON, ROY A., (10)
WINSTEAD, N. N., (272)
WOODBRIDGE, C. G., 258
YARWOOD, C. E., 85, (649)
YOUNG, H. C., Jr., (248), (444), (446)
YOUNG, ROBERT E., (102)

ZERKEL, R. S., (426)
ZILLER, WOLF G., 90, 327
ZIMMER, D. E., 944
ZUCKERMAN, BERT M., 253

SUBJECT INDEX

Abies lasiocarpa: tumors, cause unknown 473
Acer glabrum var. douglasii: Cytospora sp., pathogenicity studies, 500
--- platanoides: Cristulariella pyramidalis, 1st rept. on this host (North Carolina), 152
Aceria slykhuisii, 220
--- tulipae, 220
Achaparramiento, see under corn: stunt (virus)
Acorus calamus: Septocylindrium aromaticum, in United States, 145
Agrobacterium tumefaciens, 823
Agropyron repens: Helminthosporium leaf spot, 1st report from New Hampshire, 926; tar spot, 1st report from New Hampshire, 926
Agrostis tenuis var. Astoria: Ophiobolus patch disease in Washington, 296
Alabama, 38, 298, 517
Albizzia lebbeck: rust (Sphaerophragmium acaciae), 1st rept. in Florida, 560
Alfalfa (see also Medicago sativa): influence on soil microorganisms, 517; Verticillium albo-atrum, New Mexico biotype, 549; yellow leaf blotch, 1st rept. from New Hampshire, 926
Allium vineale: Heterosporium sp., 471
Almond, see Prunus
Alternaria sp(p.), on Arizona cypress, (secondary fungus), 98; on hop clover, 927; on Taxus, 527; in corn seed, 212; on Carthamus 146; on cotton fibers, 550; decay of Latham red raspberry, 301
--- capsule mold, of castorbean, 125
--- leaf spot, of Pieris japonica, 938; of Ricinus communis, 124
--- longipes, 159
--- ricini, 125
--- solani, 128, 539
--- tenuis, 539, 938
Amanita frostiana, 78
American beachgrass, see Ammophila
Ammophila breviligulata: Ditylenchus radicicola, 1st report in U. S. (Rhode Island) and on this host, 26
Angular leaf spot, of tobacco, 140
Annosus root rot, of slash pine plantations in Gulf States, 38
Announcements: "Index of Photographs in PHYTOPATHOLOGY, Vols. 1-41, 314; Supplement series discontinued with Supplement 262, 238
Anthracnose, of Juglans nigra, 167; of tobacco, 140
--- fruit rot, of tomato, 539
Antibiotics (see also Fungicides):
---, control of bean rust with phleomycin, 244
---, cycloheximide, in control of powdery mildew of blueberry, 368; for soil treatment to control cucumber diseases, 376; cycloheximide derivatives, for systemic control of rose powdery mildew, 794; cycloheximide derivatives, toxicity to roses, 794; cycloheximide semicarbazone derivative, 366
---, effects of aerial application on tree diseases and forest vegetation, 722
---, streptomycin, 260
---, use in non-fungistatic bacteriostatic medium for soil fungus studies, 866
Antirrhinum majus: rust, control with AMTP compounds, 756
Aphanomyces euteiches, 422, 569
--- raphani, 482
Aphelenchoides, in strawberry roots, 67
Aphelenchus avenae, 20
Aphids and the barley yellow dwarf virus in New Brunswick, 466
Aphis gossypii, 697, 877
--- spiraecola, 697
Apical necrosis, of ornamental plants caused by rapid temperature changes, 41
Aporcelaimus spp., 22
Apparatus, see techniques
Apple: blotch, control with fungicides, 441
---: fireblight, 141; control of with phenacridane chloride, 100
---: nematodes, control by soil treatment, 42
---: powdery mildew, 10; control studies, 106
---: root rot, control by soil treatment, 42

---: scab, effect of fungicides on fungus in leaf lesions, 759
---: soil treatment with nematocides and fungicides at planting time, effect on tree growth, 42
---: "star cracking" (virus), symptoms and transmission, 45
Apricot (see also Prunus): susceptibility to various pathogens, 659
--- ring pox, of plum, 796
Arceuthobium americanum, 472
Argentina, 235
Arizona, 241, 587, 826, 831, 851
Arkansas, 95, 96, 392, 690, 900, 957
Armillaria mellea, 389
Arrowweed: see Pluchea sericea
Arthrobotrys sp., 165
--- arthrobotryoides, 164
--- dactyloides, 165
Ascochyta sp., red clover, 698
Asparagus: root knot studies, 191
Aspergillus flavus, 550; in corn seed, 212
--- glaucus, in corn seed, 212
--- niger, 550; on apricot fruit, inoculation studies, 659; in corn seed, 212
--- ochraceus, in corn seed, 212
--- terreus, 550
--- versicolor, in corn seed, 212
Azalea, see Rhododendron

Bacteria, secondary, see secondary organisms
Bacterial black stalk, of tobacco, 141
--- blight, of cotton, 126
--- kernel rot, of corn in Wisconsin, 99
--- leaf spot, of Chrysanthemum morifolium, 16; of pepper, 874; of Ricinus communis, 125
--- necrosis, of Carnegiea gigantea, host range studies, 587
--- scab, of gladiolus, 344; of tomato, 481
--- spot, of peach, 100; of pepper, 120
--- wilt, of corn, 393
Bactericides (see also under control and under hosts)
---, phenacridane chloride, a broad-spectrum bactericide-fungicide, 933, for apple fireblight and peach bacterial spot control, 100
Bacterium, cause of so-called beet latent virus disease, 85
--- stewartii, 393
Banana: Pseudomonas solanacearum in Costa Rica, 435; Radopholus similis, eradication methods, 457
Barley: barley yellow dwarf (virus), epidemiology in New Brunswick, 466, influence of environment and growth substances on development of, 192; Helminthosporium sorokinianum, susceptibility of 16 varieties to, 941; loose smut, percentage yield loss as related to percentage infection, 348; root knot, varietal reaction to, 191; rust in Kansas 1959, 894; Rynchosporium scald, limiting factor in Guatemala, 227; take-all, 803
Basal stem rot, control of, 863; (footrot) of rye in Washington, 803
Bean: cucurbit latent virus infectious to, 677; Helminthosporium sorokinianum, natural infection in Michigan, 941; root rot, control with chitin 487; rust, control with AMTP compounds, 756, control with phleomycin, 244, race 33 of Uromyces phaseoli var. typica in Oregon, 388; translocation of streptomycin, 260
---, bush snap: rust, new race (34) associated with outbreak in Arkansas, 690
---, pinto: black root-rot development on, influenced by soil temperatures, 804; fusarium root-rot of, effects of soil temperature and crop residues on, 960; root-rot complex of, effects of barley residue on fungi involved, 808
Beet: "latent virus" determined to be a bacterium, host range, 85
---, sugar: Heterodera schachtii, 1st rept. of this nematode in New York, 151
Belonolaimus spp., 20, 59
--- gracilis, 21
--- longicaudatus, 21, 173, 1st rept. from Arkansas, 392
Berberis spp.: stem rust races, 453
Betula alba: Cytospora pathogenicity studies, 500
Birdsfoot trefoil, see Lotus corniculatus
Black root rot, of pinto beans, 804; of tobacco, 141
Black shank, of tobacco, 140, 968
Blackspot, of rose, 628, control of, 652
Black stem and leaf spot, complex of red clover, 698
Blast, of rice, 576
Blotch, of apple, 441
Blueberry: see Vaccinium
Blue mold, of tobacco, 140, 319
Book reviews: Are You Your Garden's Worst Pest, by Cynthia Westcott, 750; Bacterial Plant Pathogens, by C. Stapp, 394; Monograph of Monochaetia and Pestalotia, by Emil F. Guba, 666; Nematology, edited by J. N. Sasser and W. R. Jenkins, 79

Botryosphaeria sp., associated with summer cane blight of blueberry, 844
Botrytis, decay of Latham red raspberry, 301
--- cinerea, on strawberry, 818; on Trifolium spp., 927
Bougainvillea glabra: Cercospora bougainvilleae, 1st rept. on this host in El Salvador, 666
Brazil, 416
British Columbia, 90, 204, 327, 854
Bromus spp.: Helminthosporium sorokinianum, 1st report on B. brevis, B. frondosus, B. tomentosus (N.Y.), 199
--- inermis: Rhynchosporium scald, 1st report from New Hampshire, 926
--- willdenovii: head smut, control of, 216
Brown patch, of turf grasses, 870
Brown root rot, in dead jack pine, 836
Brown spot, of tobacco, 159
Brown spot disease, of Zizania aquatica, in Minnesota, 901
Brussels sprouts: fumigation for control of sugar-beet nematode, yield response to, 739
Buxus spp.: nematodes in North Carolina, 492

Cabbage: tipburn (non-parasitic), varietal reaction, 29
Cacti: Erwinia carnegieana pathogenic to, 587
Caeoma pinitorquum, 327
Caladium sp.: Meloidogyne arenaria, in Guatemala, 425
California, 89, 90, 151, 164, 232, 264, 268, 475, 562, 577, 598, 632, 636, 641, 649, 677, 682, 700, 702, 705, 709, 712, 739, 772, 796, 831, 851, 869, 940
Camellia: flower blight of, spread and distribution in Louisiana, 856; Glomerella dieback of grafts, prevention with water-emulsifiable asphalt, 432; nematodes, in North Carolina, 495
Canker (fungus undet.), of American sycamore, 78
Canna indica: rust, control with AMTP compounds, 756
Cantaloupe: cucumber mosaic virus, 137; root knot, varietal reaction to, 191; tobacco ringspot virus, 137; watermelon mosaic virus, 137
Carnegiea gigantea: bacterial necrosis, host range studies, 587
Carrot: cavity spot (physiological) of roots, secondary organisms, 102, 105
Carthamus spp. and vars.: Cercospora carthami, 1st rept. from Israel, 153, seed-borne 153; diseases in Israel, studies on susceptibility, 146; Puccinia carthami, races in Israel, 146
--- tinctorius: powdery mildew, 1st rept. in U. S. (Ariz.) 826
Carya cordiformis: Hypoxylon spp., 196
Caryopteris incana: Meloidogyne hapla control, preplant only vs. supplemental applications, 380
Castorbean: see Ricinus
Cavity spot, of carrot and parsnip roots, 102
Celery: early blight (Septoria apii), sporulation of fungus reduced by oils, 282; Fusarium yellows, control by soil fumigation, 76, importance in Colorado, 76
Cenchrus pauciflorus: wheat streak mosaic virus, 222
Cephalosporium acremonium, in corn seed, 212
Ceratocystis fagacearum, 386, 736, 749; in oak leaves, 839; spread in Missouri, importance of root grafts, 506
--- fimbriata, on apricot fruit, inoculation studies, 659
--- ulmi, 74, 141, 180, 309, 681, 749, 900; zinc chloride not effective against, 152
Cercospora, on Carthamus, 146
--- apii, 140
--- arachidicola, 674
--- bougainvilleae, 666
--- carthami, 153
--- leaf spot, of castorbean, 125, of red clover, 698
--- nicotianae, see C. apii
--- personata, 674
--- ricinella, 125
--- thujina, 96, 731, on new hosts, 745
Cercospora zebrina, 698, 927
Cereals: rusts, epidemiology in Kansas 1959, 894
Chaetomium sp., antagonistic to Fusarium, 808
Chamaecyparis pisifera: Cercospora thujina, 1st rept. (North Carolina), 745
Chauistle, Aztec name for wheat rust in Mexico, 632
Chemicals, new, for control of powdery mildew of apples, 107
Chenopodium amaranticolor: infected by cucurbit latent virus, 677
Cherry (see also Prunus): Eola rasp leaf (virus), 520; rosette, non-identity with Pfeffingerkrankheit and affinity with Stecklinburger disease, 228; tomato ring spot virus isolated from Eola rasp leaf of, 520

Chitin, and the biological control of Fusarium diseases, 487
Chrysanthemum: Stemphylium ray speck, flower rot symptoms associated with, 798
--- morifolium: bacterial leaf spot, in Florida, 16; control of Pythium root and basal stem rot with Dexon, 863
Citrus sp(p.): burrowing nematode, 22, control of, 454; citrus nematode, 23; ectoparasitic nematodes associated with decline, 23; nematodes associated with roots of, in Florida, 20; Phytophthora brown rot, contact infections prior to hot water treatment, 268, control, 264, 270; Radopholus similis, control with Cynem, 782, control with DBCP and fungicides, 62, effect on mineral (K, N, P) content of leaves, 564, gnotobiotic techniques, in study of, 742; root microflora, effect of fungicides, 65; "Sour orange seedling tree decline" in Spain, 945; tristeza virus, systemic spread in affected tree, 697, varietal reaction, effect of rootstock and scion, 416; Tylenchulus semipenetrans, in Israel, 505; vein-enation virus transmitted by Aphis gossypii, 877; wound-induced woody galls on Rough lemon and sour lime trees infected with vein-enation virus, 682
---, limon: Penicillium digitatum, biphenyl for control, 477
--- reticulata: cachexia, inverse pitting, and xyloporosis (viruses) in Israel, 356
Cladosporium sp., in corn seed, 212; on hop clover, 927; decay of Latham red raspberry, 301
Claviceps paspali, insect vectors, 530
Climates, arid, plant disease forecasting in, 31
Clover: See Trifolium
Clover: sweet, see Melilotus
Coconut: nematodes associated with yellow mottle decline in Guam, 599; yellow mottle decline (?virus) in Guam, possible spread by dagger nematodes, 411, 599
Coffea excelsa: ring spot virus of, seed transmission, 185
Coleosporium vernoniae, on pine in Arkansas, 957
Colletotrichum atramentarium, 539
--- destructivum, 140, 298
--- lagenarium, 373
--- trifolii, 299
Colombia, 334, 954
Colorado, 76, 472, 831
Common leaf spot, of red clover, 699
Conifer tumors, cause unknown, in the central Rocky Mountains, 472
Conoderus vespertinus, 225
Connecticut, 528, 720
Control (see also antibiotics, bactericides, fungicides, nematocides, etc.,) of anthracnose of walnut, 167; of bacterial diseases of fruit trees with phenacridane chloride, 100; of bacterial leaf spot of chrysanthemum, with chemical sprays, 16; of bacterial scab and Fusarium corm rot of gladiolus, with chemicals, 344; of bacterial spot of peach and fireblight of apple with phenacridane chloride, 100; of bean rust with phleomycin, 244; of blackspot of rose, 630, 652; of blotch of apple with fungicides, 441; of Botrytis rot of strawberries with fungicides, 818; of cotton seedling diseases by in-the-furrow application of soil fungicides, 276; of diseases and insect pests of crop plants in the Soviet Union, 399; of Dutch elm disease, with chemicals, 681, studies on chemical destruction of undesirable elms, 180; of Fusarium (new race) on resistant tomato varieties with chemicals, 972; of Fusarium yellows of celery, by soil fumigation, 76; of head smut in Bromus willdenovii, 216; of Heterodera schachtii, 741; of internal cork virus in sweetpotato with insecticides (negative results), 330; of leaf spot of iris, 470; of needle blight of Arizona cypress, 731; of nematodes with fertilizer-nematocide mixtures, 620, of nematode damage with potassium fertilization of infested cotton fields, 932; peanut leaf spot, 674; of pecan scab, 142; of Phytophthora brown rot of lemon with hot water treatment, 264; of plant rusts with AMTP compounds, 756; of powdery mildew of apples in Washington, 106, of blueberry, 368, of rose, 628, of roses with derivatives of cycloheximide, 794; of Pratylenchus penetrans on narcissus, 290; of Pythium root and basal stem rot of Chrysanthemum with Dexon, 863; of Pythium root diseases with fungicides, 858; of Radopholus similis on banana, 458, on citrus with Cynem, 782, with DBCP and fungicides, 62, in Citrus orchards, comparison of preplant soil treatment chemicals for, 454; of ratoon stunting disease of sugarcane by hot-water

(Control) treatments, 954; of red raspberry fruit decay with sulfur dioxide fumigation, 301; of Rhizopus fruit rot of sweet cherry by U-2069, 636; of root-knot nematode, 380, with Cynem, 335, predacious fungi not effective 164; of root rot of peas with soil fungicides, 569, of root rot and nematodes, of fruit trees by time-of-planting soil treatment, 42; of Sclerotinia rot on irrigated lettuce, studies in Washington, 51; of sclerotiniose of lettuce in Florida, 552; of Septoria leaf spot on celery, 282; of soil-borne pests of tomato with soil fumigation, 58; of soil fungi with pyridinethol derivatives plus PCNB, 426; of tobacco brown spot, with Dyrene, and economic aspects, 159; of tomato diseases (soil-borne pests) with chemicals, 58
--- biological, of Fusarium diseases, effect of chitin, 487
--- chemical, of weather fleck of tobacco, 583
--- plant disease, terminology, suggested changes, 263
--- systemic, of powdery mildew of roses, 366
---, tests, portable inoculation tower for use in, 907
Coriolus versicolor, 188
Corm rot and yellows, of gladiolus, 344
Corn: bacterial wilt in Illinois, 1961 forecast, 393; fungi isolated from unstored seed, 212; Helminthosporium spp., pathogenicity studies, 644; Helminthosporium turcicum, new type of resistance to, 780; kernel rot, new (bacterium undet.) in Wisconsin, 99; Pratylenchus zeae, 1st report in California, 940; Puccinia polysora (southern corn rust), 1st report on this host from Illinois, 236; Puccinia sorghi, 236; stalk rot, associated fungi, pathogenicity studies on, 208; stunt virus (3 types) in El Salvador, 281; sun scald of, 787
Correction, 236, 314, 394, 750
Costa Rica, 435, 980
Cotinus coggygria: Verticillium wilt in Rhode Island, 24
Cotton: Alternaria, 550; Aspergillus flavus, 550; Aspergillus niger, 550; Aspergillus terreus, 550; bacterial blight resistant and susceptible lines, Fusarium wilt resistance in, 126; boll rot, uncollapsed fibers associated with, 550; diseases, reduction in yield caused by, 1960, 219; Fusarium moniliforme, 550; Fusarium wilt resistance in bacterial blight resistant and susceptible lines, 126; nematodes, effect of potassium level on damage from, 932, nematode, reniform, spread of and crop failure from, in Louisiana, 671, nematode, root-knot resistance, associated with Fusarium wilt resistance, 127; Nigrospora sp., 550; Penicillium, 550; Rhizopus nigricans, 550; Rhizopus stolonifer, 550; seedling diseases of, control (by in-the-furrow soil fungicides), costs and benefits, 276; southwestern rust, control studies, 241
Cowpea, see Vigna
Crabapple, flowering: fireblight, 141
Cranberry: frost injury, reduction by fungicide treatments, 253
Crataegus douglasii: Cytospora sp., pathogenicity studies, 500
Criconemoides sp(p.), 20, 49
--- ornatum, 21
Cristulariella pyramidalis, 152
Cronartium commandrae, first rept. on Pinus echinata (Missouri) 576
--- comptoniae, new pine hosts in British Columbia, 854
--- fusiforme, 639
Crop plants: disease and insect pests control in the Soviet Union, 399
Crop residues, effect on Fusarium root rot of bean, 960
Crown gall disease, on fruit trees, in Israel, 823
Cucurbits: Colletotrichum lagenarium, relation of pigmentation and free amino acid content with resistance in watermelon, 373; Cylindrocarpon scoparium, 1st report on watermelon (Florida), 841; damping-off and soil rot of cucumber, effects of soil fungicide treatments, 376; Erwinia carnegieana, pathogenic to fruits of squash and honey dew, 589; fruit rot on cucumber, control with H3944, 200; latent virus, in California, 677; Mycosphaerella black rot of watermelon, temperature effect 557; Pellicularia rolfsii rot of watermelon, temperature effect 557; rind rot of cantaloup, 1st rept. in U.S. (Texas) 729; seed treatment, effect of storage on treated seeds 544; tobacco ringspot virus, on cucumber and watermelon 137; virus survey in Texas Lower Rio Grande Valley 137; viruses in melon growing areas, in Southwest 851; watermelon fruit rot, three uncommon types in Florida, 841; watermelon mosaic virus on cucumber, 137
Cupressus spp.: Cercospora thujina, 1st

(Cupressus spp.) rept. on C. lusitanica and C. macrocarpa (Georgia), 745
--- arizonica: blight, secondary fungi, 98; Cercospora thujina, cause of blight, 1st rept. on this host (Ark.), 96; Exosporium glomerulosum, 1st rept. on this host (Ark.), 96; needle blight of, in Georgia, occurrence and control, 731
Curvularia trifolii, 286, 298, on gladiolus, 512
Cylindrocladium scoparium, 1st rept. on watermelon (Fla.), 841
Cylindrosporium acori, see Septocylindrium aromaticum
Cynodon dactylon: Pratylenchus zeae, 1st rept. on this host (Calif.), 940
Cytospora sp., pathogenicity studies, 500
--- canker disease, of Italian prune, 918
--- chrysosperma, 591

Dactylaria brochopaga, 165
--- thaumasia, 164
Dactylella ellipsospora, 165
--- gephyropaga, 165
Dactylis glomerata: brown stripe, 1st rept. from New Hampshire, 926; Puccinia graminis var. avenae, 451
Dalbulus maidis, 281
Damping-off, of cucumber, 376
Decline, of pear, probable coidentity with moria disease, 912, in Washington 255, 258
--- of sour orange seedling trees in Spain, 945
Deficiency diseases: boron deficiency of Ilex aquifolium, 634
Delaware, 120, 190, 762, 874
Dianthus caryophyllus: nematodes associated with, in commercial greenhouses, 413
Didymellina macrospora, control of, 470
Dioscorea spp.: Meloidogyne incognita, main species on host in Guatemala, 981; nematodes of, in Guatemala, 425
Diospyros virginiana: Tylenchulus semipenetrans in Israel, 1st record on host, 505
Diphtherophora obesus, 21
--- perplexans, 21
Diplocarpon rosae, 628, 652
Diplodia maydis, in corn seed, 212
--- zeae, 866
Dipsacus fullonum: peach yellow bud (virus), in California, 649
Discolaimus spp., 22
Disease and insect pest control in the Soviet Union, 399
Distribution and prevalence, of races of Puccinia recondita in U.S. in 1960, 661
Ditylenchus, in strawberry roots, 67
--- radicicola, 1st rept. in Iceland 211; 1st rept. in U.S. and on new host, 26
Dollar spot, of turf grasses, 871
Downy mildew, of rice, 95; on small grains and grasses in Mississippi, 764
Dracaena sp.: leaf blight, Phyllosticta draconis, in India, 203
--- fragrans var. massangeana: apical necrosis, 41
Dutch elm disease, 141, 180, 749; control, 681; in Arkansas, 900; in North America, 74
Dwarf bunt, of wheat, 663

Early blight, of tomato, 539; of potato and tomato, 128
Echinochloa colonum: hoja blanca virus, in Colombia, host and vector studies, 949
--- crus-galli: node smut, 1st rept. from New Hampshire, 926; Pratylenchus zeae, 1st rept. on this host (Calif.), 940; wheat streak mosaic virus, 222
Eggplant: seed treatment studies with phenacridane chloride, 933
Elaeagnus angustifolia: Cytospora sp., pathogenicity studies, 500
Elm: see Ulmus
El Salvador, 281, 576, 666, 826, 900
Elymus arenarius: Ditylenchus radicicola, 1st rept. from Iceland, 211
Endogone, in turfgrasses, 869
Ephelis trinitensis, conidial stage of Balansia claviceps, 72
Epichloë typhina, 899
Epidemic, of Phragmidium rust on rose in Louisiana, 274
Epidemiology, in arid climates, 31; of barley yellow dwarf virus in New Brunswick, 466; cereal rusts in Kansas 1959, 894; peach rosette virus in Prunus angustifolia, 304; powdery mildew of potato in Utah, 355; wheat leaf rust in Kansas 1959, 894
Equipment, see techniques
Eradication, of Radopholus similis from bananas, 457
Eragrostis cilianensis: wheat streak mosaic virus, 222
Eremocarpus setigerus: cucurbit powdery

(Eremocarpus) mildew, 1st rept. on this host (Calif.), 232
Erigeron glaucus: peach yellow bud (virus), in California, 649
Erwinia amylovora, 100, 141, 730
--- aroideae, 141
--- carnegieana, host range studies, 587
Erysiphe sp., on safflower, 944
--- cichoracearum, 232, 354
--- graminis, f. sp. tritici, 846
--- polygoni, 927, on red clover, 698
Eupatorium odoratum: Pseudomonas solanacearum, 435
Euphorbia: Erwinia carnegieana pathogenic to E. valida and E. obescens, 587
--- pulcherrima: Sphaceloma poinsettiae, 1st rept. on this host from Puerto Rico, 375
Europe, 319
Exosporium glomerulosum, 96

Fading-out, of turf grasses, 872
Fairy rings, on turf grasses, 869
Fertilizer, for control of bacterial spot of pepper, 120, 874; effect on mineral content of nematode parasitized citrus trees, 564; effect on nematode damage to cotton, 932; in mixture with nematocides, effectiveness for nematode control, 620
--- injury, of tobacco, 140
Festuca spp.: influence on soil microorganisms, 517
--- rubra var. Pennlawn: Ophiobolus patch disease, 296
Ficus elastica: apical necrosis, 41
Fireblight, of apple, 100, 141; of flowering crab, 141; of pear, 141; of Stewart strain of Bartlett pear, 730
Flooding, for control of lettuce sclerotiniose, 552
Florida, 12, 16, 20, 38, 54, 58, 62, 168, 178, 307, 376, 454, 512, 552, 560, 564, 692, 697, 742, 782, 798, 841, 972
Flower blight, of camellias, 856
Fluorescent substances in fungi (Hymenomycetes), 777
Fomes annosus, 38, 306, 352, 660, 980
Forecasting of plant diseases, in an arid climate, problems of, 31
Forecasting and warning service, in the Soviet Union, 409
Fragaria spp.: Meloidogyne hapla, relative susceptibility, 308
--- chiloensis: peach yellow bud (virus), in California, 649
Frankliniella fusca, 766
--- occidentalis, 766, 772
Frogeye, of tobacco, 140
Fruit rot, of cucumber, 200; induced by fungi isolated from clovers inoculated on green tomato, 341
Fruit trees: crown gall disease in Israel, 823; soil treatment at planting time to improve growth, 42
Fumigants, soil, 54, chloropicrin for control of celery Fusarium yellows, 76, Nemex (formerly EP 136), for control of celery Fusarium yellows, 76
Fumigation, with sulfur dioxide gas, for control of red raspberry fruit decay, 301
Fungi, associated with white clover in Southeast in mid-summer, 298, with cotton seedling diseases, 276, with stalk rot of corn, 208
---, isolated from clovers, cause of fruit rot in green tomatoes, 341, isolated from unstored corn seed, 212
---, fluorescence in, 777
---, nematode-trapping, 164
---, secondary (see also secondary organisms), associated with turfgrass diseases, 871, 872, on flowers of Ladino white clover, 286
---, soil, associated with decline of mint, 1st. report in Indiana, 288, effect of various herbicides on growth in culture, 814, effect of barley residue on, in Pinto bean root-rot complex, 808, role of, in peach replant problem, 462, vermiculite media for growing, 393
Fungicide sprays, for tomatoes resistant and susceptible to fruit cracking, 539
--- and nematocide tests, 1960, results of, 236
--- tests, for control of anthracnose of walnut, 167
--- treatments, reduction of cranberry frost injury by, 253
Fungicides (see also antibiotics, nematocides, pesticides etc.):
---: alkylmercapto-tetrahydropyrimidine compounds (AMTP) as chemotherapeutic agents, for plant rusts, 756
---: as antioxidants for control of weather fleck of tobacco, 583
---: for control of bacterial spot of pepper, 120; black spot of roses, effect of surfactants and miticides on action of, 652; blast of rice, 907; blotch of apple; 441; Botrytis rot of strawberries, in Louisiana, 818; diseases of turfgrass, 892; Fusarium patch disease, of turf grasses, 112; of Fusarium wilt of tomato, new pathogenic race, 792; leaf spot of iris, 470; leaf

(Fungicides) spot of peanut, 674; of needle blight of Arizona cypress, 731; powdery mildew of apple, 106, of blueberry, 368, of roses, 628; Pythium root diseases, 858; rust on cotton, 241; scab, of apple, studies on action of, 759; scab on pecan, 142; Sclerotinia rot of lettuce, 51, 552
---: in combination with oils, for control of Septoria leaf spot on celery, 282
---: in cucumber and pea seed-treatment storage tests, 544
---:and DBCP, in control of Radopholus similis on citrus trees, 62
---: Dexon, for control of Pythium root and basal stem rot of chrysanthemum, 863
---: Dyrene, for control of brown spot of tobacco, 159; injury to tomato, 168
---: dichlone, effect on growth, yield, and disease incidence of potato and tomato, 128
---: mercury compounds, injury to narcissus bulbs, 508
---: phenacridane chloride a broad-spectrum bactericide-fungicide, 933
---: for seed treatment of rescue grass to control head smut, 216
---, soil: for control of root rot of pea, greenhouse tests, 569, of seedling diseases of cotton, 276, of soil rot and damping-off of cucumber, 376; H3944 a new broad-spectrum soil fungicide, 200
---: systemic activity of AMTP compounds, 756
---: technique for testing effects on spore germination, 390
---: use of, at planting time, effect on fruit tree growth, 42
---: U-2069 for control of Rhizopus fruit rot of sweet cherry, 636
Fungus pathogenicity tests, production of sterile onion bulbs and roots for use in, 176
Fusarium sp(p.), 102; control of, 426; on corn, 208; isolated from white clover stolons, 299; in peach replant problem, 464; on tomato, 13
--- lateritium pini, 889
--- moniliforme, 550, 871; on corn, 208; in corn seed, 212
--- nivale, 112
--- oxysporum, 298, 871
--- --- f. apii, 76
--- --- f. batatas, 562
--- --- f. conglutinans, 487
--- --- f. gladioli, 344
--- --- f. lycopersici, 272; control of new pathogenic race, 972; influence of nitrogen and light sources on cultural characteristics, 965; races, 12
--- --- narcissi, 290, 508
--- --- retusum, 13
--- patch disease, of turf, materials tested for control, 112
--- root rot, of bean, 960; of turf grasses, 869, 871
--- roseum, 298
--- solani, 299
--- --- f. phaseoli, 487, 960; Chaetomium sp. antagonistic to, 808
--- --- pisi, 422, 569, 845
--- wilt, of cotton, 126
Fusicladium effusum, 142, 911

Ganoderma curtisii, see Polyporus curtisii
Gas sampler, portable, for measuring atmospheric oxidant, 310
Georgia, 38, 46, 196, 216, 300, 304, 307, 352, 432, 660, 692, 731, 745, 976
Gibberella fujikuroi, on apricot fruit, inoculation studies, 659
--- zeae, on corn, 208; in corn seed, 212
Gibberellic acid, effect on barley yellow dwarf, 192
Gladiolus: bacterial scab, control of, 344; corm rot and yellows, control of, 344; Curvularia disease, host-parasite relationships, 512
Gloeosporium sp., associated with summer cane blight of blueberry, 844
Glomerella cingulata on camellia, 432
Gnomonia leptostyla, 167
Gram: Stemphylium leaf spot of, 1st rept. on host (India), 979
Gramineae: diseases of forage grasses, newly reported in New Hampshire, 926; diseases of turfgrasses in southern California, 869; effect of pesticides on turfgrass diseases, 892; Helminthosporium spp., pathogenicity studies, 644; Ophiobolus patch disease in western Washington, 296; Pythium ultimum and Helminthosporium, foliar pathogens, some 1st reports of, 616
Grape: leafroll (virus), effect on powdery mildew resistance, 641; powdery mildew, resistance in leafroll-affected vines, 641, as viewed in early agricultural press in California, 700
Gray leaf spot, of tomato, 539
Gray mold, control of, 818
Growth substances, effect on barley yellow dwarf, 192

Guam, 411, 599
Guatemala, 199, 227, 425, 788, 981
Gymnosporangium fuscum, in North America, description of, 90; 1st rept. U. S. (Calif.), 151
--- globosum, 351
--- juniperi-virginianae, 351
--- sabinae, see G. fuscum

Hawaii, 868
Heliconia spp.: Pseudomonas solanacearum, 435
Helicotylenchus sp., 789
--- nannus, 20
Helminthosporium sp(p.), on oats, 926; on timothy, 926; pathogenicity to species of Gramineae, 644
--- carbonum, in corn seed, 212
--- cynodontis, 870
--- maydis, in corn seed, 212
--- oryzae, on Zizania aquatica, 901
--- sativum, see H. sorokinianum
--- sorokinianum (H. sativum), 869; on barley, var. reaction to, 941; on bean, natural infection (Mich.), 941; on Bromus spp., 199; on corn (H. sativum), 208; host-range studies, 941
--- tritici-repentis, 926
--- turcicum, 780
--- vagans, 869; as foliar pathogen of Gramineae, 618
Hemicriconemoides spp., 20; wessoni, 21
Hemicycliophora spp., 20, 59
--- arenaria, 22
--- parvana, 21, 22
--- similis, 21
Herbicides, effect on soil fungi in culture, 814; turfgrass disease, control of, 892
Heterodera sp., on Dioscorea in Guatemala, 425; on tobacco, in Virginia, 812
--- schachtii, 151, 739, 826
--- tabacum, nematode resembling, in Virginia, 812
Heterosporium sp., on Allium vineale, 471
Hibiscus cannabinus: Meloidogyne incognita acrita, 1st rept. of, in Guatemala, 119
Historical notes, on plant diseases in California, 632, 700, 702, 705, 709, 712
Hoplolaimus tylenchiformis, 20, 49
Hordeum pusillum: downy mildew, 1st rept. on this host in U. S. (Mississippi), 764
Host-parasite relationships in the Curvularia disease of gladiolus, 512
Host plants: Verticillium albo-atrum, cultural type of, 548
Host range studies, of Erwinia carnegieana, 587; of Helminthosporium vagans on grasses, 616; of Lambert mottle virus of cherry, 204; of Meloidogyne incognita var. acrita, 190; of Pythium ultimum and Helminthosporium vagans on grasses, some 1st reports, 616
Hylurgopinus rufipes, 180
Hymenomycetes, fluorescent substances in extracts from, 777; spore print preservation, 78
Hypodermella arcuata, 722
Hypoxylon tinctor, on Platanus occidentalis, 196

Iceland, 211
Idaho, 289, 591, 722, 918, 920
Ilex aquifolium: boron deficiency, 634
--- cornuta: leaf spotting following use of ovex, 300
--- crenata: nematodes, in N. Car., Meloidogyne spp., 495
Illinois, 100, 180, 244, 351, 393, 557, 780, 789
India, 203, 979
Indiana, 30, 108, 212, 288, 539, 696
Indicator hosts: for necrotic ring spot and sour cherry yellow viruses, 608
Indoleacetic acid, effect on barley yellow dwarf, 192
Insect cage, glass and cellulose acetate, 824
--- transmission, of citrus vein-enation virus, 877
--- vectors, for Claviceps paspali on Paspalum dilatatum, 530
Insecticides, for control of Dutch elm disease, 681, for control of internal cork virus in sweetpotato (negative results), 330, for control of turfgrass disease, 892
---, ovex, associated with injury to Ilex cornuta, 300
Irrigation, sprinkler, effect on soil microorganism populations, 517
Iris: leaf spot, control of, 470
Israel, 31, 146, 153, 356, 505, 823
Italy, 912

Jamaica, 457
Juglans nigra: anthracnose, control studies, 167
Juncus effusus: new host for Epichloë typhina in New Hampshire, 899
Juniperus chinensis f. globosa: cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- sargenti: cedar apple rust, 1st rept. on this host (Illinois), 351; cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- horizontalis f. plumosa: cedar-apple rust, 1st rept. on this host (Illinois), 351
--- pinchotii: cedar-hawthorn rust, 1st rept. on this host (Illinois), 351

(Juniperus) sabina: Gymnosporangium fuscum, 1st rept. in N. America, 90
--- scopulorum cv. hilli: cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- --- argentea pyramidalis: cedar hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- moffeti: cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- pendula: cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
Juniperus virginiana: Cercospora thujina, 1st rept. (Georgia and South Carolina), 745; Phomopsis blight, longevity of fungus, 134, spread in transit and storage, 136, studies on seedling survival, 134
--- --- albo-spica: cedar-apple rust, 1st rept. on this host (Illinois), 351; cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- (Berg's strain): cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- cupressifolia: cedar-apple rust, 1st rept. on this host (Illinois), 351; cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- nova: cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- reptans: cedar-hawthorn rust, 1st rept. on this host (Illinois), 351
--- --- schottii: cedar-hawthorn rust, 1st rept. on this host (Illinois), 351

Kabatiella caulivora, on red clover, 698
Kansas, 309, 373, 446, 894
Kenaf, see Hibiscus
Kentucky, 140
Kernel rot, new, of corn in Wisconsin, 99
Korea, 680

Late blight, of potato, as viewed in early agricultural press in California, 705
Leaf blight, of Dracaena sp., 203
--- blotch of Bermuda grass, 870
--- curl, of peach, as viewed in early agricultural press in California, 702
--- drop, of Taxus spp., 527
--- spot, of iris, 470; peanut, 674
Lecanosticta sp., on Pinus monticola, 722
Legumes, forage: diseases newly reported in New Hampshire, 926
Lemon: see Citrus
Lepiota molybdites, on turfgrasses, 869
Leptodiscus terrestris, 298
Leptosphaeria herpotrichoides, on rye, 803
--- pratensis, 926
Lettuce: Sclerotinia rot, control on irrigated land in Washington, 51, limiting factor in Washington, 51, soil and foliar treatments for control of, 552; spotted wilt (virus), role of thrips vectors in California, 772
Life cycle, of tree pathogens, 916
Light: effect on Fusarium oxysporum f. lycopersici in culture, 965; pea wilt and root rot favored by low intensity, 574
Liquid nitrogen, for preservation of plant viruses, 755, of rust urediospores, 384
Lolium multiflorum: downy mildew, 1st report on this host in U.S. (Mississippi), 764
Longidorus elongatus, 21
Losses: from cotton diseases in 1960, 219; from cotton seedling diseases, 276; from tobacco blue mold in West Germany, 319; from tobacco brown spot in North Carolina, 159
Lotus corniculatus: Rhizoctonia crown rot, in Va., 572; root knot studies, 191
Louisiana, 274, 298, 625, 671, 818
Lupinus sp(p.): powdery mildew, 542
--- angustifolius: Stemphylium leaf spot complex of, 725, effect on seed yield, 959
Lycopersicum spp. and hybrids: Fusarium wilt, reaction to races of pathogen, 14
--- esculentum: see tomato

Macadamia integrifolia: trunk canker caused by Phytophthora cinnamomi, 868
Macrophomina phaseoli, cause of watermelon fruit rot, 841
Macrosiphum avenae, 466
Magnolia soulangeana: Verticillium wilt, 1st rept. on this genus (Indiana), 108
Maine, 717
Malus pumila: Cytospora sp., pathogenicity studies, 500
Mandarin: see Citrus reticulata
Maryland, 10, 167, 734
Massachusetts, 102, 253, 528
Medium, for culturing fungi, use of vermiculite, 393; non-fungistatic bacteriostatic, for culturing soil fungi, 866
Melampsora pinitorqua, on Pinus ponderosa (new host) in British Columbia (1st report in North America), 327;

(Melampsora) hosts and distribution, 327
Melilotus alba: Pseudopeziza leaf spot, 1st report from New Hampshire, 927; Stagonospora leaf spot, 1st rept. from New Hampshire, 926
Meloidogyne spp., 20; control tests, 54; Meloidogyne spp., reaction of tobacco breeding lines to, 692
--- acrita, 734
--- arenaria, 734; on Dioscorea in Guatemala, 425
--- hapla, 308, 534, 734; on grape-hyacinth, 746
--- incognita, 46, 577, 734; control, effect on sweetpotato yield, 497, predacious fungi not effective, 164; on Dioscorea in Guatemala, 425, 981
--- --- acrita, 46, 54, 119, 126, 976; control of 335; host range studies 191; on Tabebuia, 826
--- --- incognita, 58
--- javanica, 54, 577, 734; physiological studies on host-parasite relationship, 928
--- thamesi, 734
Melting-out, of bluegrass, 870
Mentha spp.: stolon decay in Indiana, 288
--- cardiaca: Septoria leafspot, oil yield reduced due to, epiphytotic proportions of, in northern Indiana, 696
Method (see also Technique): for adsorption and elution of tobacco mosaic virus in an ECTEOLA-cellulose column, 598; of growing sterile onion bulbs and roots for pathogenicity tests, 176; of inoculation in dwarf bunt investigations, 663; for long-term storage of sugarcane mosaic virus, 178; for rapid determination of necrotic ring spot and sour cherry yellow virus, 608; for rapid identification of the onion pink root fungus, 289; for separating biphenyl-sensitive from non-sensitive strains of Penicillium digitatum, 475
Metopolophium dirhodum, 466
Mexico, 632
Michigan, 109, 368, 392, 422, 544, 569, 699, 941
Microflora, root, effect of fungicides, 65
Microforge, electrical, aid to pure culture isolation, 312
Microsphaera alni, 368
Millet, cattail: see Pennisetum glaucum
Milo, see Sorghum vulgare
Minnesota, 260, 746, 749, 901
Mississippi, 38, 78, 530, 542, 639, 764
Missouri, 366, 506, 576, 794
Miticides, effect on action of fungicides, 652
Monstera deliciosa: apical necrosis, 41
Montana, 314, 472
Moria disease, of pear, probable coidentity with pear decline, 912
Mucor sp(p.), on corn, 208; in corn seed 212
Muscari armeniacum: root-knot nematode parasite of, first report (Minn.), 746
Mycosphaerella melonis, 557
Myrothecium roridum, 729
Myzus persicae, cucurbit latent virus, transmitted by, 677; yellows-type spinach virus, transmitted by, 720

Narcissus: basal rot, control, 508; Fusarium basal rot, effect of soil fumigants, 290; injury from mercury compounds, 508; Pratylenchus penetrans, control of, 290
Nebraska, 799
Needle blight, of Arizona cypress, 731
Nematocides (see also control, pesticides, etc., and hosts): for control of burrowing nematode 454, of Pratylenchus penetrans on narcissus 290, of root-knot nematode 380, effect of supplemental applications 380, effectiveness in muck soil 534, of sugar-beet nematode 739
---: effect on brussels sprouts yield 739; on fruit tree growth 42; on mineral content of nematode-parasitized citrus trees 564
---: Cynem, for control of Radopholus similis 782, of root knot 335
---: DBCP, for control of burrowing nematode, 62, of nematodes associated with woody ornamentals, 492, of nematodes on truck crops, studies in Georgia, 46, of Radopholus similis, in combination with fungicides, 62, of Rhizoctonia solani, 892; evaluation of formulations, application depths, root-knot control, phytotoxicity to tomatoes, 976; influence of rate and time of application to truck crops, 46; phytoxicity to truck crops in Georgia tests, 46; sprinkle irrigation method of applying, 62
---: DBCP-fertilizer mixtures, 620
---: EN 18133 for control of sting nematodes, 173
---: EP-161 and EP-162, effective in preliminary tests, 54
---: and nematocide-fertilizer mixtures, performance on vegetable crops, 620
---: polyethylene tarp for confining nema-

(Nematocides) tocides in treated soil, 534
Nematodes (see also genera, hosts, nematocides, etc.):
---: affecting vegetable crops in North Carolina, control studies, 620; associated with carnations in greenhouse, 413, with citrus roots, in Florida 20, with coconut tree disease in Guam, 411, 599, with decline of woody ornamentals in North Carolina and their control by soil treatment, 491, with strawberry soil and root samples, 67, with virus diseases, peach yellow bud, 521, yellow mottle decline virus, 411
---: effect on mineral content of citrus leaves, 564
---: gnotobiotic techniques in study of, 742
---: host-parasite relationship studies, 928
---: occurrence in Saskatchewan soils, 481
---: population variations in soybean field soil, 789, in strawberry roots, 67
---: possible vectors of coconut yellow mottle decline virus, 411
---: sample screening, substitute for Baermann funnel, 747
---: surveys for, use of aircraft in, 826
---: tolerance levels in host plants, critical method for evaluating, 930
---, burrowing: control, 62
---, grass root-gall: common name proposed for Ditylenchus radicicola, 26
---, meadow: seasonal population variations, 67
---, root-knot: on kenaf in Guatemala, 119; resistance in cotton associated with wilt resistance, 127
---, sting: control of with EN 18133, influence on yield and quality of peanuts, 173
Nematode-predacious fungi, 164
New Brunswick, 466
New disease: stunt caused by Tylenchorhynchus brevidens, of small grains (Okla.), 248
New distribution: Aphanomyces raphani in Oregon, 482; Belonolaimus longicaudatus in Arkansas, 392; Ceratocystis fagacearum on red oak in Texas, 749; Cercospora bougainvillae on Bougainvillea glabra in El Salvador, 666; Cercospora carthami on Carthamus in Israel, 153; curly-top of tomato in Costa Rica, 980; Dutch elm disease in Arkansas, 900, in Minnesota, 749; Ditylenchus radicicola in Iceland, 211, in U.S. (Rhode Island), 26; downy mildew of rice in the Western Hemisphere (Arkansas), 95; Epichloë typhina in New Hampshire, 899; Fomes root rot on Pinus spp. in Ohio, 980, on Quercus falcata and Rhus copallina in Georgia, 352; grass and legume diseases in New Hampshire, 926; Gymnosporangium fuscum, 90, 151; Heterodera schachtii on sugar beet in New York, 151; Meloidogyne incognita acrita in Guatemala, on Hibiscus cannabinus, 119; pink root of onion in Argentina, 235; Polyporus spp., on living peach trees in U.S. (S.C.), 186; potato late blight in Montana (1959), 314; powdery mildew of Psoralea tenax in United States (Mississippi), 542, of safflower in U.S., 826, 944; Pratylenchus zeae, in California, 940; Puccinia polysora on corn in Illinois, 236; Sphaceloma poinsettiae on Euphorbia pulcherrima in Puerto Rico, 375; Sphaerophragmium acaciae on Albizzia lebbeck in Florida, 560; Stemphylium leaf spot of blue lupine in Georgia and Florida, 725; wheat streak mosaic, presence in Texas, 220
New Hampshire, 128, 286, 574, 717, 845, 899, 926
New hosts: Alternaria leaf spot on Pieris japonica in Rhode Island, 938; cedar-apple and cedar-hawthorn rusts of Juniperus spp., vars. and forms (Illinois), 351; Cercospora thujina on Cupressus arizonica in Arkansas, 96, on various conifers, 745; Commandra blister rust on Pinus echinata in Missouri, 576; Cristulariella pyramidalis on Acer platanoides in North Carolina, 152; Cronartium comptoniae on Pinus muricata and Pinus radiata, 854; cucurbit powdery mildew on Eremocarpus setigerus (Calif.), 232; Cylindrocladium scoparium, on watermelon in Florida, 841; Ditylenchus radicicola on American beachgrass (Rhode Island), 26; Epichloë typhina on Juncus effusus in New Hampshire, 899; Exosporium glomerulosum on Cupressus arizonica in Arkansas, 96; Helminthosporium sorokinianum on Bromus in New York, 199; Helminthosporium vagans, 616; Melampsora pinitorqua, on Pinus ponderosa in British Columbia, 1st rept. of its occurrence in North America, 327; Myrothecium rind rot on cantaloup, in Texas, 729;

(New hosts) peach yellow bud virus, in Calif., 649; of Pratylenchus zeae in California, 940; of Pythium ultimum, 616, on peach roots (Calif.), 462; root knot nematodes on Pluchea sericea in California, 577; of Sclerophthora macrospora in Mississippi, 764; Stemphylium sarcinaeforme (new forma? on gram, in India 979; Tylenchulus semipenetrans on Diospyros virginiana in Israel, 505; Verticillium wilt on Magnolia soulangeana in Indiana, 1st rept. on this genus, 108
New Jersey, 67, 208, 263, 301, 393, 720
New Mexico, 276, 548, 549, 804, 808, 960
New York, 151, 160, 199, 228, 312, 389, 487, 528, 608, 612, 720
Nicotiana (see also tobacco): response of N. repanda, N. sylvestris, and their amphidiploid hybrid to root-knot nematodes, 734
Nigrospora sp(p.), on cotton fibers, 550; isolated from white clover stolons, 299
--- oryzae, in corn seed, 212
Nitrogen, effect of on Fusarium oxysporum f. lycopersici in culture, 965
North America, 912
North Carolina, 134, 152, 159, 173, 225, 272, 324, 330, 362, 386, 491, 620, 674, 692, 745, 889
North Dakota, 348
Northern anthracnose, of red clover, 698

Oak wilt, see Ceratocystis fagacearum
Oats: barley yellow dwarf (virus), influence of environment and growth substances on development of, 192; crown rust, reaction of supplementary differential varieties, 975, five new races of, 974; Helminthosporium leaf spot, 1st rept. from New Hampshire, 926; hoja blanca (virus), 1st rept. of natural infection (Colombia), 334; leaf rust, 1st rept. from New Hampshire, 926; redleaf (barley yellow dwarf virus), cause of defoliation in Kansas 1959, 897; root knot, varietal reaction to, 191; rusts in Kansas 1959, 894; soil-borne mosaic virus, in Washington, not transmitted by insects, 884; stem rust races, in U.S., 1960, 448
Ohio, 282, 380, 534
Oidium sp., on safflower, 826
Oils, for control of Septoria leaf spot on celery, 282
Oklahoma, 126, 142, 248, 441, 446
Okra: root knot control, predacious fungi not effective, 164
Olpidium sp., cause of stunt of small grains, 248
--- brassicae, 869
Onion: pink root, 1st rept. on this host from Argentina, 235, rapid identification, 289; "sterile" bulbs and roots for testing pathogenicity of fungus isolates, 176
Ophiobolus graminis, 803
--- --- avenae, 296
--- --- graminis, 296
Oregon, 388, 482, 520, 522, 628, 634, 891
Ornamentals, nematodes associated with decline in North Carolina, 491
Ornamental foliage plants, apical necrosis, 41
Ovacide, see Insecticides
Ozone toxicity, apparatus for measuring, 310; of tobacco, ("weather fleck"), 310, 583

Pachistima sp.: cycloheximide injury, 724
Pacific Northwest, 5
Panama, 457
Panicum capillare: wheat streak mosaic virus, 222
Paratylenchus spp., 789
Parsnip: cavity spot (physiological) of roots, secondary organisms, 102, 105
Paspalum dilatatum: insects identified as possible vectors of Claviceps paspali in Mississippi, 530
Pathogenicity, of various organisms to apricot fruit, 659
Paxillus atrotomentosus, cause of brown root of jack pine, cultural characteristics, 836
Peach (see also Prunus): bacterial spot, control of with phenacridane chloride, 100; fruit decays, 659; leaf curl, as viewed in early agricultural press in California, 702; necrotic-ring-spot virus in orchards and nursery stock, in Michigan, 109; Polyporus spp., associated with wood decay of living trees, 1st rept. on this host in U.S. (S.C.), 186; powdery mildew, 10; Pythium ultimum, 1st rept. on host (Calif.), 462; replant problem, role of fungi, 462
Peanut: leaf spot, control, 674; sting nematodes control with EN 18133, influence on yield and quality, 173
Pear: bark measles (virus-like), may be seed-borne, 891; decline (cause unknown), influence of rootstock and trends in Washington orchards, 255, 258; decline in North America, and moria disease in Italy, probable coidentity, 912; fireblight, 141, Stewart strain

(Pear) of Bartlett variety resistant to, 730; Gymnosporangium fuscum, 1st rept. from Calif., from United States (?) 151; moria, resemblance to decline, 912; quick decline (cause unknown) in Washington, 258; red leaf symptom associated with decline, 257, 259; trellis rust(Gymnosporangium fuscum) 1st rept. in North America, 90
Peas: cucurbit latent virus infectious to, 677; root rot in Michigan 422, soil fungicides for control, 569; seed treatment, effect of storage on treated seeds, 544; Thielaviopsis basicola, pathogenicity studies, 422; wilt and root rot (nonpathogenic) favored by low light intensity, 574
Pecan: scab, control with protectant fungicides, 142, occurence on Stuart variety in Georgia, 911
Pellicularia filamentosa (see also Rhizoctonia), 58, 392, 869; on apricot, inoculation studies, 659
--- praticola, 869
--- rolfsii (see also Sclerotium rolfsii), 58; cause of watermelon fruit rot, 841; on watermelon, 557
Penicillium spp., in corn seed, 212; on corn, 208; on cotton fibers, 550
--- citrinum, 65
--- digitatum, on apricot fruit, inoculation studies, 659; differential sensitivity to biphenyl among strains of, 475
Pennisetum glaucum, not attacked by inflorescence blight, 72
--- purpureum, not attacked by inflorescence blight, 72
--- --- x P. glaucum hybrid: inflorescence blight (Ephelis trinitensis), in Puerto Rico, 72
Pennsylvania, 344, 346, 616, 720
Pepper: bacterial spot, control by fertilizer and foliar sprays, 120, relation of fertility level to, 874; seed treatment studies with phenacridane chloride, 933
Peridermium harknessii, 472, 722
Peronospora tabacina, 140, in Europe, 319
Pesticides, (see also fungicides, nematocides, etc.): for control of soil-borne pests of tomato, 58
Philippines, 185
Philodendron hastatum: apical necrosis, 41
--- pertusum, see Monstera deliciosa
Phleomycin, for control of bean rust, 244
Phleum pratense: Helminthosporium leaf spot, 1st rept. from New Hampshire, 926
Phoma sp., on hop clover, 927; on red clover, 698
Phomopsis cinerascens, on apricot fruit, inoculation studies, 659
--- juniperovora, 134
--- vaccinii, 844
Phragmidium sp., on rose, outbreak in Louisiana, 274
Phycomycete, undetermined, cause of blueberry root and crown rot, 844
Phyllachora graminis, 926
Phyllosticta draconis, in India, 203
--- solitaria, 441
Physiological diseases, cavity spot of carrot and parsnip roots, 102
Phytophthora brown rot, of Citrus spp., 264, 268
--- cinnamomi, 868
--- citrophthora, 264, 268
--- fragariae, 393
--- infestans, 128, 314; early history in California, 705
--- parasitica var. nicotianae, 140, 362; virulence level of populations increased by resistant varieties, 968
--- syringae, on apricot fruit, inoculation studies, 659
Phytotoxicity, of phenylmercury sprays to white oak foliage, 346
Picea abies: Cytospora sp., pathogenicity studies, 501
Picris echioides: peach yellow bud (virus), in California, 649
Pieris japonica: Alternaria leaf spot of, 1st rept. on this host, 938
Pink root, of onion, 235, 289
Pinus spp.: differential response to chemical soil and seed treatments, 799; Fomes root rot, 1st rept. in Ohio, 980; staining-fungus root disease (root stain disease), 831; Western gall rust, control experiments with antibiotics, 722
--- banksiana: Faxillus atrotomentosus causes brown root rot in, 836
--- caribaea: Fomes annosus root rot in Gulf States, 38
--- contorta: tumors, cause unknown, 472
--- echinata: Commandra blister rust, 1st rept. on this host (Missouri), 576; infection of inoculated hybrids with southern fusiform rust, 639; root-rot (Fomes annosus) in Georgia, 352
--- elliottii var. elliottii: Fomes annosus in Georgia, 660, importance in Southeast, 306
--- monticola: blister rust control, in Idaho, 722; cycloheximide injury, 724; needle blight, control studies, 722; needle cast disease, control studies, 722

(Pinus) muricata: Cronartium comptoniae, 1st rept on this host, 854
--- ponderosa: pine twist rust (Melampsora pinitorqua) 1st rept. on this host and 1st rept. of its occurrence in North America (British Columbia), 327
--- radiata: Cronartium comptoniae, 1st report on this host, 854; pitch canker Fusarium highly pathogenic to seedlings, 889
--- strobus: Armillaria mellea, 389
--- taeda: pine needle rust, epidemics in Arkansas, 957; root-rot (Fomes annosus), in Georgia, 352
Piricularia oryzae, 576, 907
Pisum sativum: root rot caused by Fusarium solani f. pisi, 845
Pitch canker, of Pinus radiata, 889
Plant disease forecasting, in an arid climate, 31
Plantago sp.: peach yellow bud (virus), in California, 649
Platanus occidentalis: canker, decline of, in the Mississippi Delta, 78; Hypoxylon tinctor associated with canker on living trees, 196
Pluchea sericea: root-knot nematodes, 1st rept. on this host (California), 577
Plum, see Prunus
Poa pratensis: Rhizoctonia solani, control studies, 892
Podosphaera leucotricha, 10, 106
--- oxyacanthae, 10
Polyethylene film, effective for sealing fumigated soil, 58
Polyporus spp., on living peach trees, 186
Polystictus versicolor, 188
Populus sp.: Cytospora sp., pathogenicity studies, 501
Potato: dichlone treatments, effect on growth, yield, and disease incidence of, 128; early blight, control, 128; late blight, control 128, 1st rept. in Montana (1959), 314; historical note (California), 705; powdery mildew in Utah 354; seed piece treatment with chemicals, 625; Verticillium wilt, 30
Powdery mildew (see also Erysiphe, Microsphaera, Podosphaera, Sphaerotheca, Uncinula), of Carthamus spp. and vars., 146
--- --- (cucurbit), undetermined, on Eremocarpus setigerus in Calif., 232
--- --- undetermined, on Psoralea tenax and other legumes in Mississippi, 542; on safflower, 1st rept. in U.S., 826
Pratylenchus spp., 20, 789
--- brachyurus, 21, 49
--- penetrans, control, 290; in strawberry roots, 67
Pratylenchus zeae, 21, 49, 940
Prunus spp.: apricot ring pox (virus) in plum in Calif., 796; crown gall on stone fruits on almond rootstock, in Israel, 823; Cytospora sp., pathogenicity studies, 500; green ring mottle virus, virus indexing program, 687; Lambert mottle virus, host range studies, 204; necrotic ring spot and sour cherry yellows viruses, rapid determination, 608; peach yellow bud virus, endemic in native hosts in California, 649; Prunus ringspot symptoms in cherry, interrelationships with associated symptoms, in Idaho, 920, viruses associated, in Idaho, 920; Rhizopus fruit rot of sweet cherry, control with U-2069, 636; viruses of necrotic ring spot, sour cherry yellow, and green ring mottle, frequency in infected cherry orchards, 612
--- angustifolia: peach rosette virus, 304
--- armeniaca: fruit decay studies with various pathogens, 659
--- avium: Lambert mottle, symptom expression in sweet cherry varieties, 206
--- cerasus: cherry rosette virus disease, not identical with Pfeffingerkrankheit, 228, possible affinity with Stecklinburger disease, 228
--- domestica: Cytospora, effect of low temperature injury on invasion by, 591, 918
Psallioţa arvensis, on turfgrasses, 869
Pseudhalenchus anchilisposomus, 21
Pseudomonas angulata, 140
--- aptata, 85
--- calendulae, 18
--- cichorii, on chrysanthemum, 16
--- marginata, 344
--- solanacearum, in Costa Rica, 435
--- syringae, 18
--- tabaci, 140
Pseudopeziza sp., on white sweet clover, 927
--- jonesii, 926
--- trifolii, red clover, 698
Pseudotsuga menziesii: Christmas tree needle blight, control studies with antibiotics, 722; tumors, cause unknown, 472
Psoralea tenax: powdery mildew, 1st rept. on this host in United States (Mississippi), 542

Puccinia sp(p.), on turfgrasses, 869; on wheat in California, historical note, 632, early history in California, 709
--- cacabata, see P. stakmanii
--- carthami, races in Israel, 146
--- coronata, 897
--- --- avenae, new races, 974
Puccinia graminis avenae, 448, 897
--- --- tritici, 448, 633, 657, 680, 894; preservation of urediospores in liquid nitrogen, 384
--- polysora, in Illinois, 236
--- recondita, 444, 446, 661, 894, 897, 926
--- sorghi, 236
--- stakmanii, 241
--- striiformis, 778
Puerto Rico, 72, 375, 481
Pungentus spp., 22
--- textilis, 21
Puratized Agricultural Spray, 346
Pyrenochaeta terrestris, 866, in Argentina, 235; rapid identification of, 289
Pyrus communis: Cytospora sp., pathogenicity studies, 500
Pythium sp(p.), 58, 379; control of 426, 858; control with H3944, 200; isolated from white clover stolons, 299; peach replant problem, 462; on tobacco, 140, 225
--- aphanidermatum, 379, 869
--- debaryanum, 866
--- root rot, control of, 863; of turf grasses, 869
--- torulosum, 871
--- ultimum, on apricot, inoculation studies, 659; 1st report on peach (Calif.), 464; as foliar pathogen of Gramineae, 618

Quercus sp(p.): Ceratocystis fagacearum, spread through root grafts, 506; on red oak, 1st rept. in Texas, 749; symptoms in white and red oak groups, 386
--- oak wilt fungus, effect of conidial concentration on perithecial formation, 736, in leaves, 839
--- alba: Ceratocystis fagacearum, 386; phytotoxicity of phenylmercury sprays, 346
--- falcata: Fomes annosus, 1st rept. on this host from Georgia, 352
--- stellata: Ceratocystis fagacearum, 386

Races (see also strains), of Fusarium oxysporum f. lycopersici, 12, 272, 972; leaf rust, North America 1961 set of wheat varieties for leaf rust race identification, 444; of Puccinia carthami in Israel, 146; of Puccinia coronata avenae, five new, 974; Puccinia graminis in the United States in 1960, 448; of Puccinia graminis tritici, two new races in southern Korea, 680; Puccinia recondita, 444, in U. S. in 1960, 661, NA61 races in Kansas and Oklahoma, 446; of Tilletia spp., as influenced by wheat varietal populations, 5; of Uromyces phaseoli var. typica, on bean, race 33 from Oregon, 388, new race in northwest Arkansas, 690; of wheat stem rust, seedling reaction to, 657
Radish: Aphanomyces raphani, in Oregon, 1st rept. of this fungus in the Pacific Northwest, 482; vascular wilt, control with chitin, 487
Radopholus similis, 22, 62, 454, 457, 564, 742, 782
Ramularia, on Carthamus, 146
Raspberry, red: raspberry mosaic viruses in some commercial stocks in eastern U.S., 882
---, red, var. Latham: sulfur dioxide effects on reduction of post-harvest decay, 301
Reaction (see also resistance, susceptibility, etc.), of alfalfa to Verticillium albo-atrum, 549; of Citrus vars. to tristeza virus when used in various rootstocks and scion combinations, 416; of host varieties to Meloidogyne incognita acrita, 191; of rose vars. to Phragmidium rust, 274; of tomato vars. and breeding lines to Fusarium oxysporum lycopersici race 1, 272; of tomato varieties to races of Fusarium wilt fungus, 12; of wheat seedlings to new isolates of stem rust, 657; of wheat varieties to bunt, effect on racial population dynamics, 5; of Zinnia varieties to tobacco ringspot virus, 606
Resistance, of blue lupine varieties to Stemphylium botryosum and S. solani, 726; in blue lupine to Stemphylium leaf spots, 958; of corn to Helminthosporium turcicum, new type of, 780; in cowpea varieties and purelines to the cowpea yellow mosaic virus, 878; to Fusarium stem rot of sweetpotatoes, method for determining, 562; to Fusarium wilt in cotton bacterial blight resistant and susceptible lines, 126; of pear, Bartlett var. Stewart strain, to fireblight, 730; to powdery mildew in leafroll-affected grapevines, 641; of rescue grass vars. to head smut, 216;

(Resistance) of tobacco breeding lines to Meloidogyne spp., 692; of two tobacco species and their hybrids to Meloidogyne spp., 734; of tobacco varieties to black shank, effect of on build-up of virulence in fungus, 968; of watermelon to Colletotrichum lagenarium, studies on chemical factors, 373; of wheat, Suwon 92 var., to stripe rust, 778; of wheat varieties and breeding lines to powdery mildew, influenced by host developmental stage, 846
Rhabdocline pseudotsugae, 722
Rhizoctonia, sp(p.), 124; associated with mint stolon decay, 288; control of 426, control with H3944, 200; inhibitory effect on, of herbicides, 814; in peach replant problem, 462
--- solani (see also Pellicularia filamentosa), 298, 379, 392, 569, 808, 866, 870; control with H3944, 200; on Lotus corniculatus, 572; on turfgrass, 892
--- fruit rot, of tomato, 160
Rhizopus stolonifer, 550, 636
--- nigricans, 550
Rhode Island, 24, 527, 938
Rhododendron spp.: nematodes, in N. Car., 492
Rhopalosiphum maidis, 466
--- padi-fitchii, 466
Rhus copallina: Fomes annosus, 1st rept. on this host from Georgia, 352
Rhynchosporium secalis, 227, 926
Rice: blast, epidemic in El Salvador, 576, fungicide tests using portable inoculation tower, 907; downy mildew, 1st rept. on this host in Western Hemisphere (Arkansas), 95; Helminthosporium spp., pathogenicity studies, 644; hoja blanca (virus), hosts and vectors in Colombia, 949, natural occurrence in wheat and oats, 1st rept. (Colombia), 334, occurrence and vectors in El Salvador, 900
Ricinus communis: Alternaria leaf spot in Texas, 124; bacterial leaf spot, 125; capsule mold, 125; Cercospora leaf spot, 125; seedling disease (undetermined) in Texas, 124
Rind rot, of cantaloup, 728
Root and crown rot (undetermined Phycomycete), of highbush blueberry, 844
Root knot, on grape-hyacinth, 746
Root rot, of bean, 487; of fruit trees, control of by soil treatment, 42; of narcissus caused by Pratylenchus penetrans, 290; of peas, 422, 569, 845; of Pinus taeda, 352; of slash pine, 306
Root-rot complex, of Pinto bean, 808
Root stain disease, of Pinus spp., 831
Rosa sp(p.): blackspot, control studies, 628, influence of surfactants and Aramite on control of, by Phaltan, 652; Phragmidium rust epidemic in Louisiana, varietal reaction, 274; powdery mildew, control studies, 628, systemic control with cycloheximide derivatives, 794, with semicarbazone derivative of Acti-dione, 366
--- multiflora: Pratylenchus spp., in N. Car., 495
Rotylenchulus reniformis, 671
Rust (see also under various genera and hosts), of cereals in Kansas 1959, 894; control with AMTP compounds, 756; of red clover, 699; of turf grasses, 872; of wheat, as viewed in early agricultural press in California, 709
Rye: downy mildew, 1st rept. on this host (Mississippi), 764; leaf rust in Kansas 1959, 897; Leptosphaeria herpotrichoides, in Washington, 803; root knot, varietal reaction to, 191

Safflower: see Carthamus
Salix alba: Cytospora sp., pathogenicity studies, 500
Saskatchewan, 481
Scab, of apple 759; Euphorbia pulcherrima, 375; pecan 142, 911
Scald, of barley, in Guatemala, 227
Schizaphis graminum, 466
Sclerophthora macrospora, 95, 764
Sclerotinia camelliae, 856
--- homoeocarpa, 869
--- sclerotiorum, 51, 552
--- trifoliorum, red clover, 698
Sclerotium bataticola, 298; inhibitory effect of herbicides on, 814
--- cepivorum, 866
--- rolfsii (see also Pellicularia rolfsii), 517, 621; antagonistic soil microorganisms, factors affecting populations, 517; control with H3944, 200; inhibitory effect of herbicides on, 814; on apricot fruit, inoculation studies, 659; on tobacco, 225
Scolecotrichum graminis, 926
Scolytus multistriatus, 180; control, 681
Scutellonema brachyurum, 21
Secondary organisms, associated with blight of Arizona cypress, 98; associated with cavity spot of carrot and parsnip, 102, 105
Seed-borne diseases: see transmission

by seed
Seed-borne fungi, in corn seed, 212
Seed disinfestation (see also seed treatment): with phenacridane chloride, a broad-spectrum bactericide-fungicide, 933
Seed treatment (see also seed disinfestation): effect on stand and vigor of ponderosa and Austrian pine, 799; effect on yield and stands of wheat, 522; effect of storage on vigor of fungicide-treated cucumber and pea seeds, 544; for control of rescue grass head smut, 216; H3944 effective in studies with vegetable crops, 200; of potato seed pieces, studies on effectiveness and safety of chemicals, 625
Seedling disease, of cotton, control of, 276; of Ricinus communis, 124
Septocylindrium aromaticum, 145
Septoria apii, 282; on apricot fruit, inoculation studies, 659
--- leafspot, of Scotch spearmint, 696
--- tritici, 788
Setaria viridis: wheat streak mosaic virus, 222
Small grains: stunt caused by Olpidium sp., 248; stunt caused by Tylenchorhynchus brevidens, a new disease, (Okla.), 248
Smoke bush: see Cotinus
Smut, of wheat, as viewed in early agricultural press in California, 712
Sodium arsenite, effective in Dutch elm disease control, 180
Sogata cubana, 949
--- orizicola, 334, 900, 949
Soil fumigants: effect on sweetpotato yields, 497; EP-161 and EP-162, preliminary report on nematocidal effectiveness, 54
Soil fumigation (see also nematocides, etc.): for control of celery Fusarium yellows, 76, of soil-borne fungi, nematodes, weeds, in tomato plantings, 58; effect on incidence of Fusarium basal rot of narcissus, 290, on stand and vigor of ponderosa and Austrian pine, 799; polyethylene film an effective seal for treated areas, 58
Soil microorganisms: influence of crop and sprinkler irrigation on populations, 517
Soil rot, of cucumber, 376
Soil temperature, effect on Fusarium root rot of bean, 960
Soil treatment (see also control, fungicides, soil fumigation, etc.): for control of sclerotiniose of lettuce, 552, of soil rot and damping-off of cucumber, 376, of tomato diseases, 58; for fruit trees, 42; use of polyethylene film in, 58; with pyridinethiol derivatives plus PCNB, effectiveness against soil fungi, 426
Soil-borne fungi, affecting tomato, control by soil fumigation, 58; control studies, 426; medium for culturing, 866
--- --- pathogens, H3944 a broad-spectrum fungicide for control of, 200
--- --- pests, of tomato, control of, 58
Solanum aviculare: Meloidogyne arenaria, in Guatemala, 425
Sorbus sitchensis: Cytospora sp., pathogenicity studies, 500
Sorghum: Helminthosporium spp., pathogenicity studies, 644
--- vulgare: Pratylenchus zeae, on milo, 1st rept. in California, 940
South Carolina, 186, 200, 298, 306, 692, 745
Soviet Union, 399
Soybean: nematodes, seasonal variations in populations, 789; powdery mildew, 542; root knot, varietal reaction to, 191
Spain, 945
Sphaceloma poinsettiae, 375
Sphaerophragmium acaciae, 560
Sphaerotheca pannosa, 10, 366, 628, 794
Spinach, cucurbit latent virus infectious to, 677; yellows-type virus, aphid-transmitted, in eastern United States, 720
Spore prints, preservation of, 78
Squash: see under cucurbits
Stachys bullata: peach yellow bud (virus) in California, 649
Stalk rot, of corn, studies on associated fungi, 208
"Star cracking", of apple, in Washington, 45
Stem rots, in tobacco, relation to wireworm injury, 225
Stem rust, of wheat, 680
Stemphylium on Carthamus, 146
--- botryosum, 725, 959
--- floridanum, 798
--- leaf spot, of gram, 979
--- ray speck, of chrysanthemum, 798
--- sarcinaeforme, 698, 979
--- solani, 539, 959
Stolon decay, of Mentha spp., 288
Storage, of fungicide-treated pea and cucumber seeds, effect on seed vigor, 544

Strains (see also races): of Penicillium digitatum, differential sensitivity to biphenyl among, 475
---, of Phytophthora parasitica var. nicotianne, 362; effect of varietal resistance on virulence level, 968
---, of Pseudomonas solanacearum from virgin soils in Costa Rica, 435
---, of Stemphylium botryosum on blue lupine, 725
Strawberry (see also Fragaria): Botrytis rot, chemical control of, 818; nematodes in root and soil samples in New Jersey, 67; Pratylenchus penetrans, seasonal population variations, 67
Streptomycin, translocation in bean, 260
Stripe rust, 778
Stunt, caused by Tylenchorhynchus brevidens, a new disease of small grains, 248
Sugar-beet nematode, survey, (aerial), 826
Sugarcane: mosaic virus, long term storage of, 178; ratoon stunting disease (virus) of, in Colombia, 954
Sulfur dioxide, fumigation with, for control of postharvest decay of Latham red raspberries, 301
Sun scald, of corn, 787
Surfactants, effect on fungicide action, 652
Survey, aerial, for sugar-beet nematode, 826
Susceptibility, of apricot fruits to various pathogens, 659; of barley varieties to Helminthosporium sorokinianum, 941; of Fragaria spp. to Meloidogyne hapla, 308; of pine hybrids to fusiform rust, 639; of Pinus radiata to pitch canker, 889; of safflower varieties and species to foliage diseases in Israel, 146; of Stuart pecans to scab, 911
Sweetpotato: Fusarium stem rot, method for determining resistance to, 562; internal cork virus, dissemination not controlled with insecticides, 330; root knot, control, 497; soil fumigation, effect on production, 497

Tabebuia pentaphylla: root-knot nematode, in El Salvador, 826
Taphrina deformans, early history in California, 702
Taxus spp.: leaf drop associated with Alternaria, 527
Technique (see also Method): conidiation test, dependability and sensitivity in detecting viable spores of oak wilt fungus, 736; for determining the preservation of plant viruses in liquid nitrogen, 755; electrical microforge, aid to pure culture isolation, 312; for evaluating tolerant levels in nematized host plants, 930; gnotobiotic, definition of term, 742, use in study of Radopholus similis on citrus, 742; for handling thrips in transmission experiments with the tomato spotted wilt virus, 766; inoculation tower, portable, for use in fungicide tests, 907; liquid nitrogen for preservation of plant viruses, 755; substitute for a Baermann funnel for screening samples of nematodes, 747; for testing effects of soluble postharvest fungicides on spore germination, 390
Techniques and apparatus: gas sampler for measuring atmospheric oxidant, 310; insect cage of glass and cellulose, 824
Temperature, rapid changes in, cause of apical necrosis of ornamental foliage plants, 41; soil, effect on black root rot of pinto beans, 804
Tennessee, 386
Terminology, of plant disease control, suggested changes, 263
Texas, 124, 138, 220, 749, 851
Thielaviopsis basicola, 141, 569, 804, 808, pathogenicity of, on peas, 422
Thrips tabaci, 766, 772
Thuja orientalis: Cercospora thujina, 96
Tilletia sp., on wheat, early history in California, 710
--- caries, racial population dynamics, 5
--- contraversa, 663
--- foetida, racial population dynamics, 5
Tipburn, of cabbage varieties, 29
Tobacco (see also Nicotiana): angular leaf spot, 140; anthracnose, 140; bacterial black stalk, 141; black root rot, 141; black shank, influence of resistant varieties on virulence of fungus, 968, spread in Kentucky, 140; blue mold, 140, in Europe, 319; brown spot (Alternaria) control, 159; cold injury, 140; cyst nematode, resembling Heterodera tabacum, in Virginia, 812; etch, aphid-transmitted virus, 140; fertilizer injury, 140; frogeye, 140; manganese injury, 140; mosaic virus, adsorption and elution of, in an ECTEOLA-cellulose column, 598; Phytophthora parasitica var. nicotianae, influence of host passage on virulence, 362; Pythium spp., 225; Pythium soft rot, 140; root-knot nematodes, reaction of breeding lines to, 692, root-knot nematodes, reaction of two Nicotiana spp. and their amphidiploid hybrid to, 734; Sclerotium rolfsii, 225; stem rot incidence in relation to wireworm injury, 225; weather fleck,

(Tobacco) (ozone toxicity), 310, chemical control, 583; wildfire, 140
Tomato: anthracnose fruit rot control, 539; bacterial scab, 1st rept. in Puerto Rico, 481; curly-top virus disease, 1st rept. in Costa Rica, 980; DBCP toxicity to, 46; dichlone treatments, effect on growth, yield and disease incidence of, 128; Dyrene phytotoxicity effects on, 168; early blight control, 128, 539; Erwinia carnegieana, pathogenic to fruits of, 589; fruit crack rot development in susceptible and resistant varieties, delayed harvest fungicide test, 539; Fusarium oxysporum lycopersici, race 1, reaction of breeding lines and vars. to, 272; Fusarium wilt, infection of resistant varieties by new strain in Florida, 12; gray leaf spot control, 539; green fruits, as medium for inducing fruit rot and sporulation with fungi isolated from clovers, 341; phytotoxicity of DBCP, effect of formulations and application depths, 976; seed treatment studies with phenacridane chloride, 933; spotted wilt (virus), role of thrips vectors in California, 772; Rhizoctonia fruit rot, effect on mold count, 160, symptoms and studies, 160; Rhizoctonia stem canker, 392; root knot, control with DBCP, 976; root knot control, predacious fungi not effective, 164; root knot, varietal reaction to, 191; Sclerotium rolfsii control with H3944, 200; soil-borne pests, soil treatment for control of, 58; wilt, chemical control of new race of fungus, 972, effect of light and nitrogen sources on fungus in culture, 965
Toxicity, manganese, to tobacco, 140
Tribulus terrestris: Pratylenchus zeae, 1st rept. on this host (Calif.), 940
Trichoderma sp., on corn, 208
--- lignorum, 299
Trichodorus spp., 20
--- aequalis, 21
--- christiei, 21, 49, 59
--- porosus, 21
--- proximus, 21
Trifolium spp.: fungi isolated from, induced rotting in green tomatoes, 341; phyllody, potato culture associated with high incidence of, in Maine and New Hampshire, 717
--- ---, hop clover: Alternaria sp., 927; Botrytis leaf spot, 1st rept. from New Hampshire, 927; Cercospora leaf spot, 1st rept. from New Hampshire, 927; Cladosporium sp., 927; Phoma sp., 927; powdery mildew, 1st rept. from New Hampshire, 927
--- arvense: Botrytis leaf spot, 1st rept. from New Hampshire, 927; Cercospora leaf spot, 1st rept. from New Hampshire, 927
--- hybridum: Cercospora leaf spot, 1st rept. from New Hampshire, 927
--- pratense: diseases found in Michigan, 698; influence on soil microorganisms, 517
--- repens: Botrytis leaf spot, 1st rept. from New Hampshire, 927; fungi associated with, in Southeast in midsummer, 298; influence on soil microorganisms, 517
--- --- var. Ladino: Curvularia trifolii on flowers, 1st. published record (N.H.), 286; secondary fungi on flowers, 286
Translocation, of streptomycin in bean, 260
Transmission, of Cercospora carthami, by seed, 153; of citrus vein-enation virus by Aphis gossypii, 877; of coffee ring spot virus, by seed, 185; of cucurbit latent virus by the aphid Myzus persicae, 677; of hoja blanca of rice by Sogata spp., 949; of pear bark measles, by seed, evidence for, 891; of yellows-type spinach virus by the aphid Myzus persicae, 720
Tree diseases: control experiments with antibiotics, 722
Tree pathogens, generalized life cycle, 916
Trunk canker, of Macadamia, 868
Tumors, on coniferous species, 472
Turf grasses (see also Gramineae): fungicidal tests for control of Fusarium patch disease, 112
Tylenchorhynchus spp., 20, 789
--- acti, 21
--- brevidens, cause of stunt of small grains (Okla.), 248; host-range studies, 251
Tylenchulus semipenetrans, 20; on American persimmon in Israel, 1st record on host, 505
Tylenchus, in strawberry roots, 67

Ulmus spp.: Dutch elm disease, 141; control studies, 180; distribution in North America, 74; dormant sprays for control of vector, 681; 1st rept. in Arkansas, 900; spread in Kansas, 309; zinc chloride not effective against fungus, 152

(Ulmus) americana: Dutch elm disease, distribution in North America, 1959, 74, 1st rept. in Minnesota, 749
Uncinula necator, leafroll-affected grape, resistant to, 641; as viewed in early agricultural press in California, 700
Uromyces phaseoli var. typica, 244, 388; new race (34) in Arkansas, 690
--- trifolii, red clover, 698
Ustilago bullata, 216
--- crus-galli, 926
--- nuda, 348
Utah, 354, 663

Vaccinium sp.: cycloheximide injury, 724; powdery mildew control, 368; root and crown rot (undetermined Phycomycete) of highbush blueberry in New Jersey, 844; summer cane blight, associated with low temperature injury and fungi, 844
Vascular wilt, of radish, 487
Vectors (see also transmission), of hoja blanca virus 949; of viruses, techniques for handling in transmission tests, 766; of wheat streak mosaic virus in Texas, 220
Vegetable crops: H3944 as a seed protectant, 200; nematode control, with DBCP, effectiveness and toxicity, 46, with EP-161 and EP-162, experimental soil fumigants, 54, with nematocides and nematocide-fertilizer mixtures, studies on performance, 620
Venturia inaequalis, 759
Vermiculite medium, for growing fungi, 393
Vernonia sp.: Coleosporium vernoniae, associated with pine needle rust in Arkansas, 957
Verticillium albo-atrum, 24, 30, 549; cultural type of, in New Mexico, 548; on Magnolia, 108
Verticicladiella sp., on Pinus spp., 831
Vigna sinensis: "beet latent virus" a bacterium, 85; cowpea yellow mosaic virus, sources of resistance to, 878
Virginia, 572, 720, 761, 812
Virus diseases (see also under hosts):
---: apricot ring pox of plum, 796
---: barley yellow dwarf virus, and aphids, in New Brunswick, 466; influence of environment and growth substances on severity, 192
---: beet latent virus (so called) is a bacterium, 85
---: cachexia and xyloporosis of mandarins, 356
---: cadang-cadang, see yellow mottle decline
---: cherry rosette, non-identity with Pfeffingerkrankheit and affinity with Stecklinburger disease, 228
---: confusion of bacteria and viruses, 85
---: cowpea yellow mosaic virus, varietal reaction to, 878
---: cucumber mosaic in cucurbits, 137, 851
---: of cucurbits, in the lower Rio Grande Valley of Texas, 137, in Southwest melon-growing areas, 851
---: curly-top of tomato in Costa Rica, 980
---: etch (aphid-transmitted) of tobacco, 140
---: green ring mottle virus of Prunus spp., 687; frequency in infected trees, 612
---: hoja blanca, grass hosts in Colombia, 949; of Echinochloa colonum, 949; of rice, 949, occurrence and insect vectors in El Salvador, 900; of wheat and oats, natural infection in Colombia, 334
---: internal cork of sweetpotato, 330
---: Lambert mottle virus of cherry, host-range studies, 204
---: latent virus in cucurbits, 677; transmitted by Myzus persicae, 677; host-range studies, 677
---: leafroll of grapes, effect on powdery mildew resistance, 641
---: mosaic of sugarcane, long term storage of virus, 178
---: necrotic ring spot in cherry, frequency in infected orchard trees, 612, rapid determination of virus, 608; in peach orchards and nursery stock, 109
---: nematodes as possible vectors, of coconut disease in Guam, 411
---: peach rosette in Prunus angustifolia, epidemiology, 304
---: peach yellow bud virus, endemic in California, 649; hosts in California, 649; relation to tomato ring spot virus, 650
---: Pfeffingerkrankheit not identical with cherry rosette, 228
---: phyllody in Trifolium spp., association with potato culture, 717
---: plum line-pattern virus in cherry, not commonly associated with ringspot symptoms in Idaho, 920
---: prune dwarf virus, association with chlorotic line symptoms in cherry in Idaho, 920
---: Prunus ringspot symptoms in cherry,

(Virus) associated viruses in Idaho, 920
---: Prunus virus indexing program ineffective when partial, 687
---: raspberry mosaic viruses in red raspberry, 882
---: ratoon stunting disease, of sugarcane, in Colombia, 954
---: ring spot of Coffea excelsa, seed transmission, 185
---: soil-borne mosaic of oats in Washington, 884; host-range studies, 886
---: sour cherry yellows virus, frequency in infected trees, 612, rapid determination of, 608
---: sour orange seedling tree decline (?virus), in Spain, 945
---: spinach yellows, transmitted by Myzus persicae, 720
---: squash mosaic virus in cucurbits in Southwest, 851; in squash, 137
---: "star cracking" of apple in Washington, 45
---: Stecklinburger disease, relationship with cherry rosette, 228
---: stunt (achaparramiento) of corn, in El Salvador, 281
---: tobacco mosaic, adsorption and elution of, in an ECTEOLA-cellulose column, 598
---: tobacco ring spot, in cucurbits, 137, 851; Zinnia varieties, reaction, 605
---: tomato ring spot virus, isolated from Eola rasp leaf of cherry, 520; relation to peach yellow bud virus, 650
---: tomato spotted wilt virus, new convolvulaceous hosts, 775; role of thrips vectors in California, 772; thrips-handling techniques in transmission experiments, 766
---: tristeza, in Citrus varieties, 416; spread of virus in affected tree, 697
---: vein-enation virus of citrus 682; Aphis gossypii a vector of, 877; role in wound-induced woody galls, 682
---: virus preservation with liquid nitrogen, 755
---: watermelon mosaic in cucurbits, 137, 851
---: wheat-streak mosaic in wheat and grasses in Texas, 220
---: xyloporosis, and cachexia, of mandarin, 356
---: yellow mottle decline of coconut (? virus), in Guam, 411, 599; possibly caused by soil-borne virus spread by dagger nematodes, 411
Washington, 41, 45, 106, 112, 255, 258, 290, 296, 508, 687, 730, 778, 803, 884
Water-emulsifiable asphalt plus fungicides to protect camellia grafts, 432
Weather, and plant diseases in arid climates, 31
"Weather fleck", of tobacco, see ozone toxicity
Weather injuries: frost injury to cranberries, 253; low temperature injury a factor in Cytospora invasion, 591; low temperature injury associated with summer cane blight of blueberry, 844; sun scald of corn due to low temperatures followed by heavy dew, 787
Weeds: control with soil treatment, 55, 60
West Virginia, 470
Wheat: bunt, influence of host varietal reaction on races of Tilletia spp., 5; races of, in Pacific Northwest, 5; as viewed in early agricultural press in California, 712
---: dwarf bunt, effect of straw cover and method of inoculation, 663
---: hoja blanca (virus), 1st rept. of natural infection (Colombia), 334
--- leaf rust, epiphytotic in Kansas 1959, 894; races, supplemental (NA61) differential varieties, 444; races in U.S. in 1960, 661
---: powdery mildew, early record in California, 710; resistance to infection by Erysiphe graminis f. sp. tritici, influenced by the stage of development of host plant, 846
---: root knot, varietal reaction to, 191
---: rust, control with AMTP compounds, 756; during Spanish era in California, 632; as viewed in early agricultural press in California, 709
--- seed-treatment fungicides, effect on yield and stands, 522
---: Septoria leaf blotch in Guatemala, 788
---: stem rust, isolates of, seedling reaction to, 657; in Kansas, epidemiology in 1959, 894; races in Korea, 680, races in U.S. in 1960, 448
---: stripe rust, not observed in Kansas 1959, 894; resistance of Suwon 92 to, 778
---: stunt in Oklahoma, 248
---: take-all, 803
---: wheat-streak mosaic in Texas, 220
"White spots", of Carthamus spp. and vars. 146
Wildfire, of tobacco, 140
Wilt and root rot (nonpathogenic), of peas, 574
Wisconsin, 29, 99, 152, 192, 681, 787, 826,

(Wisconsin) 836
Woody galls, of Citrus spp., associated with vein-enation virus infection, 682
Wyoming, 472

Xanthomonas pruni, 100
--- ricinicola, 125
--- vesicatoria, 120, 481, 874
Xiphinema spp., 20
--- americanum, 20; occurrence in Saskatchewan soils, 481; and X. diversicaudatum, associated with yellow mottle decline of coconut in Guam, 411
--- campinense, 21
--- chambersi, 21
--- diversicaudatum, 411

Zinnia: tobacco ringspot virus, varietal reaction studies, 605
Zizania aquatica: brown spot disease, epidemic in Minnesota, 901

ERRATA

On page 65, last line in 5th paragraph, read Penicillium citrinum instead of Penicillium cintrinum.

On page 560, read Albizzia lebbeck (L.) Benth. instead of Albizia lebbeck (L.) Benth.

On page 617, in Table 1, read Echinochloa crusgalli var. frumentacea instead of Echinochloa crusgalli frumantacea.

See also corrections on pages 236, 314, 394, 750.

CROPS RESEARCH DIVISION